



行政院國家科學委員會專題研究計劃 RRPA89050032 (1).P

* 在 Measure Chain 中非線性微分方程式之環上正解的存在性 *

* Positive Solutions in an Annulus for Nonlinear *

* Differential Equations on a Measure Chain *

* *

計劃類別：☒ 個別計劃 ☐ 整合型計劃

計劃編號：NSC89-2115-M-032-004

執行期間：88 年 8 月 1 日至 89 年 7 月 31 日

個別型計劃：計劃主持人：錢傳仁
國科會助理：黃纓茹
國科會助理：羅慧君
共同主持人：

整合型計劃：總計劃主持人：
子計劃主持人：

註：整合型計劃總報告與子計劃成果報告請分開編印各成一冊
，彙整一起繳送國科會。

處理方式：☒ 可立即對外提供參考

(請打√) ☐ 一年後可對外提供參考

☐ 兩年後可對外提供參考

(必要時，本會得展延發表時限)

執行單位：淡江大學數學系
中華民國 89 年 12 月 14 日

摘要

本文主要在證明一個 measure chain 上的二階微分方程式，

$$u^{\Delta\Delta}(t) + f(u(\sigma(t))) = 0, \quad t \in [0, 1],$$

滿足邊界條件

$$\alpha y(0) - \beta y^{\Delta}(0) = 0 \quad \gamma y(\alpha(1)) + \delta y^{\Delta}(\sigma(1)) = 0,$$

具有至少兩個解，而且它的解分佈解空間中的兩個環上。其中 f 為一連續正函數，且

$$(i) f_0 = \infty \quad \text{and} \quad f_{\infty} = \infty, \quad \text{or} \quad (ii) f_0 = 0 \quad \text{and} \quad f_{\infty} = \infty, \quad \text{where,}$$

f 滿足

$$f_0 = \lim_{x \rightarrow 0^+} \frac{f(x)}{x} \quad \text{and} \quad f_{\infty} = \lim_{x \rightarrow \infty} \frac{f(x)}{x},$$

方法是應用巴那赫空間上的錐形定點定理 (kransnosel'skii's 定點定理) 得證兩個解。

關鍵詞：微分方程式、特徵值、邊界值問題、measure chain。

POSITIVE SOLUTIONS IN AN ANNULUS FOR NONLINEAR DIFFERENTIAL EQUATIONS ON A MEASURE CHAIN

CHUAN JEN CHYAN, JOHNNY HENDERSON AND HUI CHUN LO

Abstract. We study the existence of positive solutions of the second order differential equation in an annulus on a measure chain, $u^{\Delta\Delta}(t) + f(u(\sigma(t))) = 0$, $t \in [0, 1]$, satisfying the boundary conditions, $\alpha y(0) - \beta y^{\Delta}(0) = 0$ and $\gamma y(\sigma(1)) + \delta y^{\Delta}(\sigma(1)) = 0$, where f is a positive function and $f(x)$ is sublinear (respectively superlinear) at $x = 0$ and is superlinear (respectively sublinear) at $x = \infty$. The methods involve applications of a fixed point theorem for operators on a cone in a Banach space.

1. Introduction

In this paper, we study the existence of positive solutions of a certain second order differential equation in an annulus on a measure chain. Much recent attention has been given to differential equations on measure chains, and we refer the reader to [4], [9] and [15] for some historical works as well as to the more recent papers [1], [1], [2], [5], [6], [10], [11] and the book [19] for excellent references on these types of equations. Before introducing the problems of interest for this paper, we present some definitions and notation which are common to the recent literature. Our sources for this background material are the two papers by Erbe and Peterson [10], [11].

Definitions 1.1. Let T be a closed subset of \mathbb{R} , and let T have the subspace topology inherited from the Euclidean topology on \mathbb{R} . For $t < \sup T$ and $r > \inf T$, define the *forward jump operator*, σ , and the *backward jump operator*, ρ , respectively, by

$$\begin{aligned}\sigma(t) &= \inf\{\tau \in T \mid \tau > t\} \in T, \\ \rho(r) &= \sup\{\tau \in T \mid \tau < r\} \in T,\end{aligned}$$

for all $t, r \in T$. If $\sigma(t) > t$, t is said to be right scattered, and if $\rho(r) < r$, r is said to be left scattered. If $\sigma(t) = t$, t is said to be right dense, and if $\rho(r) = r$, r is said to be left dense.

Received March 22, 1999; revised May 14, 1999.

1991 *Mathematics Subject Classification.* 34B99, 39A99

Key words and phrases. Differential equation, measure chain, boundary value problem.

Definitions 1.2. For $x : T \rightarrow R$ and $t \in T$ (if $t = \sup T$, assume t is not left scattered), define the *delta derivative* of $x(t)$, $x^\Delta(t)$, to be the number (when it exists), with the property that, for any $\epsilon > 0$, there is a neighborhood, U , of t such that

$$\left| [x(\sigma(t)) - x(s)] - x^\Delta(t)[\sigma(t) - s] \right| \leq \epsilon |\sigma(t) - s|,$$

for all $s \in U$. The second delta derivative of $x(t)$ is defined by $x^{\Delta\Delta}(t) = (x^\Delta)^\Delta(t)$.

If $F^\Delta(t) = h(t)$, then define the *integral* by

$$\int_a^t h(s) \Delta s = F(t) - F(a).$$

Throughout this paper, we assume T is a closed subset of R with $0, 1 \in T$.

Definition 1.3. Define the closed interval, $[0, 1]$, in T by

$$[0, 1] = \{t \in T \mid 0 \leq t \leq 1\}.$$

Other closed, open, and half-open intervals in T are similarly defined.

In this paper, we consider the existence of positive solutions of the differential equation on a measure chain,

$$u^{\Delta\Delta}(t) + f(u(\sigma(t))) = 0, \quad t \in [0, 1], \quad (1)$$

satisfying

$$\alpha y(0) - \beta y^\Delta(0) = 0 \quad \gamma y(\sigma(1)) + \delta y^\Delta(\sigma(1)) = 0, \quad (2)$$

where $f : R^+ \rightarrow R^+$ is continuous and $d = \gamma\beta + \alpha\delta + \alpha\gamma(\sigma(1)) > 0$.

We remark that by a solution, u , of (1), (2), we mean $u : [0, \sigma^2(1)] \rightarrow R$, u satisfies (1) on $[0, 1]$, and u satisfies the boundary conditions (2).

This paper constitutes an extension of the recent work by Erbe and Peterson [11] in which they obtained positive solutions of both (1), (2) for the cases when either $f(x)$ is superlinear (at $x = 0$ and $x = \infty$), or $f(x)$ is sublinear (at $x = 0$ and $x = \infty$). The solutions obtained in [11] were found to belong to the intersection of a cone with an annular type region. In a more recent paper, Erbe and Peterson [12] applied index theory in a cone to obtain multiple positive solutions of (1), (2). Our goal is to extend to differential equations on measure chains the works of Erbe, Hu, and Wang [13], Erbe and Tang [14], and Kaufmann [17], to obtain two positive solutions of (1), (2), when f is superlinear at one endpoint (zero or infinity) and sublinear at the other (respectively, infinity or zero); that is when either (i) $f_0 = \infty$ and $f_\infty = \infty$, or (ii) $f_0 = 0$ and $f_\infty = 0$, where,

$$f_0 = \lim_{x \rightarrow 0^+} \frac{f(x)}{x} \quad \text{and} \quad f_\infty = \lim_{x \rightarrow +\infty} \frac{f(x)}{x}.$$

We will make our assumptions on f more precise in Section 3. The results herein are also related to those by Atici [3] and Eloe and Henderson [8].

A Krasnosel'skii fixed point theorem [20] is applied to yield positive solutions of (1), (2) in our arguments. These methods have been used successfully, in the cases when (1) is either an ordinary differential equation or a finite difference equation; see [17], [18], [22].

In Section 2, we present some properties of a Green's function associated with (1), (2) that are used in defining a positive operator. We also state the Krasnosel'skii fixed point theorem. In Section 3, we give an appropriate Banach space and construct a cone on which we apply the fixed point theorem yielding twin positive solutions of (1), (2).

2. Some Preliminaries

In this section, we state the above mentioned Krasnosel'skii fixed point theorem. We will apply this fixed point theorem in the next section to a completely continuous integral operator whose kernel, $G(t, s)$, is the Green's function for

$$-y^{\Delta\Delta}(t) = 0, \quad (3)$$

$$\alpha y(0) - \beta y^{\Delta}(0) = 0 \quad \gamma y(\sigma(1)) + \delta y^{\Delta}(\sigma(1)) = 0. \quad (4)$$

Erbe and Peterson [11] have found

$$G(t, s) = \begin{cases} \frac{1}{d} \{ \alpha t + \beta \} \{ \gamma(\sigma(1) - \sigma(s)) + \delta \}, & t \leq s, \\ \frac{1}{d} \{ \alpha \sigma(s) + \beta \} \{ \gamma(\sigma(1) - t) + \delta \}, & \sigma(s) \leq t, \end{cases} \quad (5)$$

where

$$d = \gamma\beta + \alpha\delta + \alpha\gamma(\sigma(1)) > 0,$$

from which

$$G(t, s) > 0, \quad (t, s) \in (0, \sigma(1)) \times (0, 1), \quad (6)$$

$$\frac{G(t, s)}{G(\sigma(s), s)} = \begin{cases} \frac{\alpha t + \beta}{\alpha \sigma(s) + \beta}, & t \leq s, \\ \frac{\gamma(\sigma(1) - t) + \delta}{\gamma(\sigma(1) - \sigma(s)) + \delta}, & \sigma(s) \leq t. \end{cases} \quad (7)$$

This implies

$$G(t, s) \leq G(\sigma(s), s) \quad \text{for } t \in [0, \sigma(1)], \quad s \in [0, 1],$$

and it is also shown in [11] that

$$G(t, s) \geq kG(\sigma(s), s), \quad t \in \left[\frac{\sigma(1)}{4}, \frac{3\sigma(1)}{4} \right], \quad s \in [0, 1], \quad (8)$$

where $k = \min \left\{ \frac{\alpha\sigma(1)+4\beta}{4\alpha\sigma(1)+4\beta}, \frac{\gamma\sigma(1)+4\delta}{4\gamma(\sigma(1)-\sigma(0))+4\delta} \right\}$.

We will apply the following fixed point theorem which can be found in the book by Krasnosel'skii [20] as well as in the book by Deimling [7].

Theorem 2.1. *Let B be a Banach space, and let $P \subset B$ be a cone in B . Assume Ω_1, Ω_2 are open subsets of B with $0 \in \Omega_1 \subset \overline{\Omega}_1 \subset \Omega_2$, and let*

$$T : P \cap (\overline{\Omega}_2 \setminus \Omega_1) \rightarrow P$$

be a completely continuous operator such that either,

- (i) $\|Tu\| \leq \|u\|$, $u \in P \cap \partial\Omega_1$, and $\|Tu\| \geq \|u\|$, $u \in P \cap \partial\Omega_2$, or
 (ii) $\|Tu\| \geq \|u\|$, $u \in P \cap \partial\Omega_1$, and $\|Tu\| \leq \|u\|$, $u \in P \cap \partial\Omega_2$.

Then T has a fixed point in $P \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

3. Solutions of (1.1), (1.2) in a Cone

In this section, we apply Theorem 2.1 to the eigenvalue problem (1), (2). Throughout this section, we assume $\sigma(1)$ is right dense so that $G(t, s) \geq 0$, for $t \in [0, \sigma^2(1)]$, $s \in [0, \sigma(1)]$.

Assume also throughout this section that $[0, \sigma(1)]$ is such that

$$\xi = \min\{t \in T \mid t \geq \frac{\sigma(1)}{4}\}, \text{ and}$$

$$\omega = \max\{t \in T \mid t \leq \frac{3\sigma(1)}{4}\}$$

both exist and satisfy,

$$\frac{\sigma(1)}{4} \leq \xi < \omega \leq \frac{3\sigma(1)}{4},$$

and if $\sigma(\omega) = 1$, also assume $\sigma(\omega) < \sigma(1)$. Next, let $\tau \in [\xi, \omega]$ be defined by

$$\int_{\xi}^{\omega} G(\tau, s) \Delta s = \max_{t \in [\xi, \omega]} \int_{\xi}^{\omega} G(t, s) \Delta s. \quad (9)$$

Finally, we define

$$\ell = \min_{s \in [0, \sigma(1)]} \frac{G(\sigma(\omega), s)}{G(\sigma(s), s)}, \quad (10)$$

and set

$$m = \min\{k, \ell\}. \quad (11)$$

We note that $u(t)$ is a solution of (1), (2), if and only if,

$$u(t) = \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s, \quad t \in [0, \sigma^2(1)].$$

For our constructions, let $B = \{x : [0, \sigma^2(1)] \rightarrow R \mid x \text{ is bounded}\}$, with norm $\|x\| = \sup\{|x(t)| : t \in [0, \sigma^2(1)]\}$. Then, define the cone $P \subset B$

$$P = \{x \in B \mid x(t) \geq 0 \text{ on } [0, \sigma^2(1)], \text{ and } x(t) \geq m\|x\|, \text{ for } t \in [\xi, \sigma(\omega)]\}.$$

Theorem 3.1. Assume that f satisfies the following conditions:

- (A) $f_0 = \infty$ and $f_{\infty} = \infty$.

(B) *There exists a $p > 0$ such that, if $0 \leq x \leq p$, then,*

$$f(x) \leq \eta p$$

where,

$$\eta = \left(\int_0^{\sigma(1)} G(\sigma(s), s) \Delta s \right)^{-1}$$

Then there exist at least two solutions u_1, u_2 of (1), (2) which belong to P , such that

$$0 \leq \|u_1\| \leq p \leq \|u_2\|.$$

Proof. Define an integral operator $T : P \rightarrow B$ by

$$Tu(t) = \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s, \quad u \in P. \quad (12)$$

We seek a fixed point of T which belongs to P , and so we first claim $T : P \rightarrow P$. We choose $u \in P$. Then from (7),

$$\begin{aligned} Tu(t) &= \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s \\ &\leq \int_0^{\sigma(1)} G(\sigma(s), s) f(u(\sigma(s))) \Delta s, \end{aligned}$$

and so

$$\|Tu\| \leq \int_0^{\sigma(1)} G(\sigma(s), s) f(u(\sigma(s))) \Delta s. \quad (13)$$

So, from (8) and (13), for $u \in P$,

$$\begin{aligned} \min_{t \in [\xi, \omega]} Tu(t) &= \min_{t \in [\xi, \omega]} \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s \\ &\geq \int_0^{\sigma(1)} k G(\sigma(s), s) f(u(\sigma(s))) \Delta s \\ &\geq k \|Tu\| \\ &\geq m \|Tu\|, \end{aligned}$$

and from (10)

$$\begin{aligned} Tu(\sigma(\omega)) &= \int_0^{\sigma(1)} G(\sigma(\omega), s) f(u(\sigma(s))) \Delta s \\ &\geq \int_0^{\sigma(1)} \ell G(\sigma(s), s) f(u(\sigma(s))) \Delta s \\ &\geq m \int_0^{\sigma(1)} G(\sigma(s), s) f(u(\sigma(s))) \Delta s \\ &\geq m \|Tu\|. \end{aligned}$$

Thus $Tu \in P$, and we conclude $T : P \rightarrow P$. The standard arguments show T is completely continuous.

From $f_0 = \infty$ there is an $p > r > 0$ such that $f(u) \geq Mu$, for $0 \leq u \leq r$, where the constant $M > 0$ satisfies

$$Mm \int_{\xi}^{\omega} G(\tau, s) \Delta s \geq 1. \quad (14)$$

Let

$$\Omega_1 = \{x \in B \mid \|x\| < r\},$$

and then choose $u \in P$ with $\|u\| = r$. Then

$$\begin{aligned} Tu(\tau) &= \int_0^{\sigma(1)} G(\tau, s) f(u(\sigma(s))) \Delta s \\ &\geq \int_{\xi}^{\omega} G(\tau, s) f(u(\sigma(s))) \Delta s \\ &\geq M \int_{\xi}^{\omega} G(\tau, s) u(\sigma(s)) \Delta s \\ &\geq Mm \int_{\xi}^{\omega} G(\tau, s) \Delta s \|u\| \\ &\geq \|u\|. \end{aligned}$$

Thus, $\|Tu\| \geq \|u\|$, and in particular,

$$\|Tu\| \geq \|u\|, \text{ for } u \in P \cap \partial\Omega_1. \quad (15)$$

Now consider $u \in P$ with $\|u\| = p$. From condition (B),

$$\begin{aligned} Tu(t) &= \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s \\ &\leq \int_0^{\sigma(1)} G(\sigma(s), s) f(u(\sigma(s))) \Delta s \\ &\leq \int_0^{\sigma(1)} G(\sigma(s), s) \Delta s p \eta \\ &= \|u\|. \end{aligned}$$

If we define $\Omega_2 = \{u \in B : \|u\| < p\}$, then

$$\|Tu\| \leq \|u\|, \quad u \in P \cap \partial\Omega_2. \quad (16)$$

Theorem 2.1, together with (15) and (16), implies that there exists a fixed point, u_1 , satisfying $r < \|u_1\| < p$.

Next, using condition (A) again, we know there exists an $H_1 > 0$ such that

$$f(u) \geq \varepsilon u, \quad (17)$$

for all $u \geq H_1$, where we choose $\varepsilon > 0$ such that,

$$\varepsilon m \int_{\xi}^{\omega} G(\tau, s) \Delta s \geq 1. \quad (18)$$

Let $H = \max\{2p, \frac{H_1}{m}\}$ and pick $u \in P$ with $\|u\| = H$. Since $u(t) \geq m\|u\| \geq H_1$ on $t \in [\xi, \sigma(\omega)]$, we have

$$\begin{aligned} Tu(\tau) &= \int_0^{\sigma(1)} G(\tau, s) f(u(\sigma(s))) \Delta s \\ &\geq \varepsilon \int_{\xi}^{\omega} G(\tau, s) u(\sigma(s)) \Delta s \\ &\geq \varepsilon m \int_{\xi}^{\omega} G(\tau, s) \Delta s \|u\| \\ &\geq \|u\|. \end{aligned}$$

Set $\Omega_3 = \{u \in B : \|u\| < H\}$. Then

$$\|Tu\| \geq \|u\|, \quad u \in P \cap \partial\Omega_3. \quad (19)$$

Theorem 2.1, together with (16) and (19), implies that there exists a fixed point, u_2 , of T such that $p < \|u_2\| < H$ and the proof is complete.

We now consider the case $f_0 = 0$ and $f_{\infty} = 0$.

Theorem 3.2. Assume that f satisfies the following conditions:

(A') $f_0 = 0$ and $f_{\infty} = 0$.

(B') There exists a $q > 0$ such that, if $mq \leq x \leq q$, then,

$$f(x) \geq \lambda q$$

where,

$$\lambda = \left(\int_{\xi}^{\omega} G(\tau, s) \Delta s \right)^{-1}.$$

Then there exist at least two solutions u_1, u_2 of (1), (2) which belong to P , such that

$$0 \leq \|u_1\| \leq q \leq \|u_2\|.$$

Proof. From $f_0 = 0$ there is an $0 < r < q$ such that $f(u) \leq \eta u$ for $0 \leq u \leq r$ where the constant

$$\eta = \left(\int_0^{\sigma(1)} G(\sigma(s), s) \Delta s \right)^{-1}.$$

Let

$$\Omega_1 = \{x \in B \mid \|x\| < r\}.$$

Then

$$\begin{aligned} Tu(t) &= \int_0^{\sigma(1)} G(t, s) f(u(\sigma(s))) \Delta s \\ &\leq \int_0^{\sigma(1)} G(\sigma(s), s) \eta u(\sigma(s)) \Delta s \\ &\leq \eta \int_0^{\sigma(1)} G(\sigma(s), s) \Delta s \|u\| \\ &\leq \|u\|. \end{aligned}$$

If we define $\Omega_1 = \{u \in B : \|u\| < r\}$, then

$$\|Tu\| \leq \|u\|, \quad u \in P \cap \partial\Omega_1. \quad (20)$$

Now consider $u \in P$ with $\|u\| = q$. Then, from (B') ,

$$\begin{aligned} Tu(\tau) &= \int_0^{\sigma(1)} G(\tau, s) f(u(\sigma(s))) \Delta s \\ &\geq \int_{\xi}^{\omega} G(\tau, s) f(u(\sigma(s))) \Delta s \\ &\geq \lambda \int_{\xi}^{\omega} G(\tau, s) \Delta s q \\ &\geq \|u\|. \end{aligned}$$

Thus, $\|Tu\| \geq \|u\|$, and in particular, if we define $\Omega_2 = \{u \in B : \|u\| < q\}$, then

$$\|Tu\| \geq \|u\|, \quad \text{for } u \in P \cap \partial\Omega_2. \quad (21)$$

Theorem 2.1, together with (20) and (21), implies that there exists a fixed point, u_1 , satisfying $r < \|u_1\| < q$. Returning to condition (A') , we know that for any $\varepsilon > 0$, there exists an $M > 0$ such that,

$$f(u) \leq M + \varepsilon u \quad \text{for } u \geq 0, \quad 0 \leq t \leq 1.$$

And so,

$$\begin{aligned} Tu(t) &\leq \int_0^{\sigma(1)} G(t, s) [M + \varepsilon u(\sigma(s))] \Delta s \\ &\leq M \int_0^{\sigma(1)} G(\sigma(s), s) \Delta s + \varepsilon \int_0^{\sigma(1)} G(\sigma(s), s) u(\sigma(s)) \Delta s. \end{aligned}$$

We choose $\varepsilon > 0$ sufficiently small and $L > \frac{M}{\eta}$ sufficiently large, so that for $u \in P \cap \partial\Omega_3$

$$\|Tu\| \leq L = \|u\|, \quad (22)$$

where

$$\Omega_3 = \{x \in B : \|u\| < L\}.$$

We obtain the desired result $0 \leq \|u_1\| \leq q \leq \|u_2\|$ by appealing to (20), (21), (22) and Theorem 2.1. This completes the proof.

Acknowledgement

This paper is supported by the National Science Council of the Republic of China.

References

- [1] R. P. Agarwal and M. Bohner, Basic calculus on time scales and some of its applications, preprint.
- [2] R. P. Agarwal, M. Bohner and P. Wong, *Sturm-Liouville eigenvalue problems on time scales*, Appl. Math. Comput., in press.
- [3] F. Atici, *Two positive solutions of a boundary value problem for difference equations*, J. Difference Eqns. Appl. 1(1995), 263-270.
- [4] B. Aulback and S. Hilger, *Linear dynamic processes with inhomogeneous time scale*, Nonlinear Dynamics and Quantum Dynamical Systems, Math. Res. 59, Akademie Verlag, Berlin, 1990.
- [5] C. J. Chyan and J. Henderson, *Eigenvalue problems for nonlinear differential equations on a measure chain*, J. Math. Anal. Appl., in press.
- [6] C. J. Chyan, J. M. Davis, J. Henderson and W. K. C. Yin, *Eigenvalue comparisons for nonlinear differential equations on a measure chain*, Electronic Journal of Differential Equations 1998(1998), 1-7.
- [7] K. Deimling, Nonlinear Functional Analysis, Springer-Verlag, New York, 1985.
- [8] P. W. Elloe and J. Henderson, *Positive solutions and nonlinear $(k, n-k)$ conjugate eigenvalue problems*, Differential Equations Dynam. Systems 6(1998), 309-317.
- [9] L. H. Erbe and S. Hilger, *Sturmian theory on measure chains*, Differential Equations Dynam. Systems 1(1993), 223-246.
- [10] L. H. Erbe and A. Peterson, *Green's functions and comparison theorems for differential equations on measure chains*, Dynam. Contin. Discrete Impuls. Systems, in press.
- [11] L. H. Erbe and A. Peterson, *Positive solutions for a nonlinear differential equation on a measure chain*, Math. Comput. Modelling, in press.
- [12] L. H. Erbe and A. Peterson, *Eigenvalue conditions and positive solutions*, preprint.
- [13] L. H. Erbe, S. Hu and H. Wang, *Multiple positive solutions of some boundary value problems*, J. Math. Anal. Appl. 184(1994), 640-648.
- [14] L. H. Erbe and M. Tang, *Existence and multiplicity of positive solutions to nonlinear boundary value problems*, Differential Equations Dynam. Systems 4(1996), 313-320.
- [15] S. Hilger, *Analysis on measure chains - a unified approach to continuous and discrete calculus*, Resultate Math. 18(1990), 18-56.