

Contents

Chapter 0 Introduction	1
Chapter 1 Linear Density Response	3
1.1 Non-Relativistic Hamiltonian	3
1.1.1 The Lagrangian for many-body systems	3
1.1.2 Canonical momentum	3
1.1.3 Non-relativistic Hamiltonian	4
1.1.4 Quantization of the electromagnetic field	4
1.1.5 Intrinsic spin in the magnetic field	5
1.1.6 Born-Oppenheimer approximation	5
1.1.7 Model Hamiltonian for electrons	6
1.2 Non-Resonant Inelastic Photon Scattering	7
1.2.1 Time dependent perturbation theory (TDPT)	7
1.2.2 Inelastic photon scattering	8
1.3 Linear Density Response	12
1.3.1 Density-density response function	12
1.3.2 Excitation energy of N-particle system	14
1.3.3 Connection with dynamic structure factor	16
1.4 Time-Dependent Density-Functional Theory	18
1.4.1 Density-functional theory (DFT)	18
1.4.2 Kohn-Sham response function	21
1.4.3 Meaning of Kohn-Sham response function	23
1.4.4 Application to phonon calculation	25
Chapter 2 Excitons in NiO and CoO	28
2.1 Introduction	28
2.2 Geometric structure	30
2.3 Electronic structure	31
2.3.1 LDA+U method	31
2.3.2 Band structure	33

- with
lent
68
- Fig. 2.72 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Hartree-Fock case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for CoO. 81
- Fig. 2.73 q -dependence of the imaginary parts of χ for the Hartree-Fock case in CoO with the energies at (a) 1.37eV, (b) 2.32eV, (c) 2.81eV, and (d) 3.17eV. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 82
- Fig. 3.1 Cubic structure of PbTiO_3 . 86
- Fig. 3.2 Tetragonal structure of PbTiO_3 . 86
- Fig. 3.3 Tetragonal structure of $\text{Pb}_{0.75}\text{Ca}_{0.25}\text{TiO}_3$. 87
- Fig. 3.4 Primitive unit cell of cubic $\text{Pb}_{0.5}\text{Ca}_{0.5}\text{TiO}_3$. 87
- Fig. 3.5 Phonon dispersion curves of tetragonal PTO corresponding to Type I and Type II calculations (see in the text) are shown as solid and dotted curves, respectively. Open symbols (circles, triangles)[85] and solid symbols (circles[71], diamonds[83], squares[84]) denote various experimental data. 91
- Fig. 3.6 Sound velocities of longitudinal (L) and two shear waves (S1 and S2) of tetragonal PTO in (a) a-b and (b) a-c planes. Shear wave polarizes nearly perpendicular (parallel) to the plane is represented as S1 (S2). Brillouin scattering experiment data[72] and calculations based on experimental and theoretical relaxed geometries are plotted as dashed, solid and dotted lines, respectively. 93
- Fig. 3.7 TO1 mode of cubic PbTiO_3 with frequency $215i \text{ cm}^{-1}$. The direction of wave vector q is pointed out on the right-hand side of the plot. 97

2.3.3 Wannier function	42
2.4 Excitons	45
2.4.1 Experimental results	45
2.4.2 Property of excitons	48
2.4.3 Angular dependence	50
2.4.4 Energy of exciton in NiO	54
2.4.5 Energy of exciton in CoO	69
2.5 Summary	83
Chapter 3 Phonons in PbTiO₃ and Related Mixing Material	84
3.1 Introduction	84
3.2 Geometric structure	86
3.3 Computational method	88
3.4 Results and Discussion	89
3.4.1 Structural detail	89
3.4.2 Lattice dynamics of tetragonal PbTiO ₃	90
3.4.3 Dynamical effective charge analysis of PbTiO ₃	93
3.4.4 Doping effect in Pb _{1-x} Ca _x TiO ₃	103
3.4.5 Phase transitions in Pb _{1-x} Ca _x TiO ₃	105
3.5 Summary	107
Chapter 4 Conclusion	108
Bibliography	110

List of Tables

- 42
- 45 **Table 2.1** Fractional coordinates of selected \mathbf{k} points for band-structure plots of NiO and
45 CoO 42
- 48
- 50 **Table 3.1** Calculated (Type I, Type II (see in the text) and GGA[82]) and observed
54 (Expt.[82]) lattice constants (in a.u.), c/a ratios, and internal parameters (in
69 fractional coordinates along the z direction (u)) for tetragonal PTO. O1, O2 and
83 O3 denotes the oxygen atom on the yz , xz , and xy face of the unit cell,
84 respectively. 90
- 84
- 86 **Table 3.2** Optimized internal parameters for tetragonal $\text{Pb}_{0.75}\text{Ca}_{0.25}\text{TiO}_3$ corresponding to
88 experimental lattice constants. O1, O2 and O3 denotes the oxygen atom on the
89 yz , xz , and xy face of the unit cell, respectively. 90
- 89
- 90 **Table 3.3** Elastic constants $c_{\mu\nu}$ in GPa; piezoelectric constants e_{iv} in C/m^2 , and
93 dielectric constants ε_{ij} (with phonon contribution) for tetragonal PTO.
03 Corresponding calculated data with respect to theoretical optimized lattice
05 constants are presented in parentheses. 92
- 07
- 08 **Table 3.4** Born effective charges, the shortest Ti-O bondlength $d_{\text{Ti-O}}$ (in a.u.) and the
0 enthalpy difference (in meV) with respect to the cubic (C) phase for the
tetragonal (T) phase of tetragonal PTO at different lattice volumes. The
calculated value of hydrostatic pressure corresponding to each smaller cell
volume is also given for reference. 95

Table 3.5 Correlation matrix and mode effective charges for cubic and tetragonal PTO at experimental lattice constants. The data in parentheses for cubic and tetragonal phases are obtained from the literature[80] and theoretical optimized lattice constants, respectively. 96

Table 3.6 Dielectric constant along c-axis ($\epsilon_{33}(\infty)$) and Born effective charges (Z_{\parallel}^*) for $\text{Pb}_{1-x}\text{Ca}_x\text{TiO}_3$ and a fictitious tetragonal CaTiO_3 (see in the text). All data are obtained at experimental lattice constants. Effective charges Z_{\parallel}^* refer to atom displacements parallel to c-axis. The oxygen at the bottom of TiO_5 in tetragonal PTO is represented as O3 and the others are denoted by O1. 96

List of Figures

- Fig. 1.1** Kohn-Sham response function $\chi_{\sigma\sigma}^{KS}(\bar{x}, \bar{x}'; \omega)$, $\chi_{\sigma\neq\sigma'}^{KS} = 0$. 21
- Fig. 1.2** Connection between many-body and Kohn-Sham response functions. 24
- Fig. 2.1** Rhombohedral structure of NiO (CoO). Anti-ferromagnetic order is constructed from (111) planes composed of Ni (Co) atoms along the [111] direction in rocksalt structure denoted schematically by small arrows. 30
- Fig. 2.2** LDA+U band structure of NiO. The bands with *s*-orbital character of (a) spin-up Ni1 (b) spin-down Ni1 (c) spin-up Ni2 (d) spin-down Ni2 are highlighted by open circles. 34
- Fig. 2.3** LDA+U band structure of NiO. The bands with *p*-orbital character of (a) spin-up Ni1 (b) spin-down Ni1 (c) spin-up Ni2 (d) spin-down Ni2 are highlighted by open circles. 35
- Fig. 2.4** LDA+U band structure of NiO. The bands with *d*-orbital character of (a) spin-up Ni1 (b) spin-down Ni1 (c) spin-up Ni2 (d) spin-down Ni2 are highlighted by open circles. 36
- Fig. 2.5** LDA+U band structure of NiO. The bands with *s*-orbital character of (a) spin-up O (b) spin-down O and *p*-orbital character of (c) spin-up O (d) spin-down O are highlighted by open circles. 37
- Fig. 2.6** LDA+U band structure of CoO. The bands with *s*-orbital character of (a) spin-up Co1 (b) spin-down Co1 (c) spin-up Co2 (d) spin-down Co2 are highlighted by open circles. 38

- Fig. 2.7** LDA+U band structure of CoO. The bands with *p*-orbital character of (a) spin-up Co1 (b) spin-down Co1 (c) spin-up Co2 (d) spin-down Co2 are highlighted by open circles. 39
- Fig. 2.8** LDA+U band structure of CoO. The bands with *d*-orbital character of (a) spin-up Co1 (b) spin-down Co1 (c) spin-up Co2 (d) spin-down Co2 are highlighted by open circles. 40
- Fig. 2.9** LDA+U band structure of CoO. The bands with *s*-orbital character of (a) spin-up O (b) spin-down O and *p*-orbital character of (c) spin-up O (d) spin-down O are highlighted by open circles. 41
- Fig. 2.10** Selected **k** paths for band-structure plots of NiO. 42
- Fig. 2.11** Energy levels of *d*-character Wannier functions in NiO and CoO. 43
- Fig. 2.12** Comparison between LDA+U within RPA and non-resonant inelastic X-ray scattering experiment for NiO at small momentum transfer ($q \sim 0.7 \text{ \AA}^{-1}$) at absolute unit. The low energy part in panel (a) is enlarged in panel (b). 45
- Fig. 2.13** Dynamical structure factors found by non-resonant inelastic X-ray scattering experiment for NiO and CoO. High-resolution (0.3 eV) measurements of (a) the *q*-dependence of the *d-d* peak intensities for NiO in the [111] direction; (b) the *q*-orientation dependence in CoO at $q = 7 \text{ \AA}^{-1}$. Panel (c) and panel (d) represent the *q*-orientation dependence for $q = 3.5 \text{ \AA}^{-1}$ in NiO and $q = 3.75 \text{ \AA}^{-1}$ in CoO, respectively. 46

- spin-up
hted by
39
- spin-up
nted by
40
- n-up O
O are
41
- 42
- 43
- X-ray
olute
45
- ering
) the
the
t the
oO,
46
- Fig. 2.14** Low-resolution (1.1 eV) non-resonant inelastic X-ray scattering measurements and LDA+U/RPA calculations of the dynamical structure factor for NiO and CoO: (a) measurements along the [100] and [111] directions for NiO; (b) similar measurements on CoO; (c) calculations for NiO along the [100] and [111] directions; (d) similar calculations for CoO. 47
- Fig. 2.15** Density fluctuation propagator in the many-body electron system. 48
- Fig. 2.16** Bethe-Salpeter equation. 49
- Fig. 2.17** A two-band ladder example for the Bethe-Salpeter equation. 49
- Fig. 2.18** (a) q -dependence of transition probability of e'_g to e_g for NiO (3.5\AA^{-1} at white line). Averaging over cubic equivalent anti-ferromagnetic domain is imposed. (b) and (c) are the polar plots of the calculated and measured d - d spectral weights, respectively. 50
- Fig. 2.19** q -dependence of transition probability of e'_g to e_g for pure cubic symmetry. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 51
- Fig. 2.20** q -dependence of transition probability of a_g to e_g for NiO. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 51
- Fig. 2.21** (a) q -dependence of transition probability of e'_g to e_g for CoO (3.5\AA^{-1} at white line). Averaging over cubic equivalent anti-ferromagnetic domain is imposed. (b) and (c) are the polar plots of the calculated and measured d - d spectral weights, respectively. 52

- Fig. 2.22** q -dependence of transition probability of e'_g to a_g for CoO. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 52
- Fig. 2.23** Wannier functions of e_g states in (a) NiO and (b) CoO. Note the bulge distortion for CoO (at arrow) compared to the nearly cubic-symmetry shape for NiO. 53
- Fig. 2.24** Wannier functions of e'_g states in (a) NiO and (b) CoO. Note the slightly less nodal shape for CoO (at arrow) compared to the nearly cubic-symmetry shape for NiO. 53
- Fig. 2.25** Illustration of (a) Fock and (b) Hartree interaction diagrams in Bethe-Salpeter equation. 55
- Fig. 2.26- 2.27** The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the L_0 case in NiO. The corresponding diagram is presented inside the plot. 56- 57
- Fig. 2.28- 2.30** The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Local case in NiO. The corresponding diagram is presented inside the plot. 57- 58
- Fig. 2.31- 2.35** The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Fock case in NiO. The corresponding diagram is presented inside the plot. 59- 61
- Fig. 2.36- 2.40** The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Hartree-Fock case in NiO. The corresponding diagram is presented inside the plot. 61- 63

Fig. 2.41 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the L_0 case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for NiO. The zero intensity along $\langle 100 \rangle$ direction reflects the nodal direction. 65

Fig. 2.42 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Local case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for NiO. The zero intensity along $\langle 100 \rangle$ direction reflects the nodal direction. 65

Fig. 2.43 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Fock case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for NiO. The zero intensity along $\langle 100 \rangle$ direction reflects the nodal direction. 66

Fig. 2.44 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Hartree-Fock case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for NiO. The zero intensity along $\langle 100 \rangle$ direction reflects the nodal direction. 66

Fig. 2.45 q -dependence of the imaginary parts of χ for the L_0 case in NiO with the energies at (a) 6.81eV and (b) 7.74eV. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 67

Fig. 2.46 q -dependence of the imaginary parts of χ for the Local case in NiO with the energies at (a) 1.65eV and (b) 1.95eV. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 67

Fig. 2.47 q -dependence of the imaginary parts of χ for the Fock case in NiO with the energies at (a) 1.43eV and (b) 1.95eV. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 68

Fig. 2.48 q -dependence of the imaginary parts of χ for the Hartree-Fock case in NiO with the energies at (a) 1.72eV and (b) 2.61eV. Averaging over cubic equivalent anti-ferromagnetic domain is imposed. 68

Fig. 2.49- 2.50 The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the L_0 case in CoO. The corresponding diagram is presented inside the plot. 70

Fig. 2.51- 2.53 The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Local case in CoO. The corresponding diagram is presented inside the plot. 71- 72

Fig. 2.54- 2.61 The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Fock case in CoO. The corresponding diagram is presented inside the plot. 72- 76

Fig. 2.62- 2.68 The real (dashed line) and imaginary (solid line) parts of polarization propagator L for the Hartree-Fock case in CoO. The corresponding diagram is presented inside the plot. 76- 79

Fig. 2.69 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the L_0 case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for CoO. 80

Fig. 2.70 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Local case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for CoO. 80

Fig. 2.71 $\chi(q \sim 7\text{\AA}^{-1}, q \sim 7\text{\AA}^{-1}; \omega)$, the results of the Fock case weighting by the Fourier transforms of Wannier functions at $q \sim 7\text{\AA}^{-1}$ for CoO. 81

- Fig. 3.8** LO3 mode of cubic PbTiO_3 with frequency 600.9 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 97
- Fig. 3.9** TO2 mode of cubic PbTiO_3 with frequency 63.0 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 98
- Fig. 3.10** LO1 mode of cubic PbTiO_3 with frequency 62.7 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 98
- Fig. 3.11** TO3 mode of cubic PbTiO_3 with frequency 442.5 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 99
- Fig. 3.12** LO2 mode of cubic PbTiO_3 with frequency 414.1 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 99
- Fig. 3.13** TO1 mode of tetragonal PbTiO_3 with frequency 145.8 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 100
- Fig. 3.14** LO1 mode of tetragonal PbTiO_3 with frequency 188.7 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 100
- Fig. 3.15** TO2 mode of tetragonal PbTiO_3 with frequency 360.2 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 101
- Fig. 3.16** LO2 mode of tetragonal PbTiO_3 with frequency 443.2 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 101
- Fig. 3.17** TO3 mode of tetragonal PbTiO_3 with frequency 646.4 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 102

f wave
97

Fig. 3.18 LO3 mode of tetragonal PbTiO_3 with frequency 784.8 cm^{-1} . The direction of wave vector \mathbf{q} is pointed out on the right-hand side of the plot. 102

vector
98

Fig. 3.19 Difference between square of $A_1(3\text{LO})$ and $A_1(3\text{TO})$ phonon frequencies for tetragonal $\text{Pb}_{1-x}\text{Ca}_x\text{TiO}_3$ from Raman experiment[68]. 103

wave
98

Fig. 3.20 Eigendisplacement of the softest mode in cubic $\text{Pb}_{0.5}\text{Ca}_{0.5}\text{TiO}_3$. 106

wave
99

wave
99

wave
00

wave
00

wave
01

wave
01