

Table of Contents

	Page
LIST OF FIGURES	v
LIST OF TABLES	vii
ACKNOWLEDGMENTS	viii
CURRICULUM VITAE	ix
ABSTRACT OF THE DISSERTATION	x
Introduction	1
1 Soliton Hierarchies from Splittings of Lie Algebra	6
1.1 Soliton Hierarchy constructed from $\mathcal{L}_{\pm}^{\tau}(\mathcal{G})$ and $\{a\lambda^j \mid j \geq 1\}$	6
1.2 Soliton Hierarchy constructed from $\mathcal{L}_{\pm}^{\tau, \sigma}(\mathcal{G})$ and $\{a\lambda^{2j} \mid j \geq 1\}$	19
1.3 Soliton Hierarchy Constructed from $\mathcal{L}_{\pm}^{\tau, \sigma}(\mathcal{G})$ and $\{a\lambda^{2j+1} \mid j \geq 1\}$	22
2 Schrödinger Flows on Hermitian Symmetric Spaces	27
2.1 Schrödinger Flows on Compact Kahler Manifolds	27
2.2 Schrödinger Flows on Hermitian Symmetric Spaces $\frac{U}{K}$ and Relation with $\frac{U}{K}$ -NLS Equations	29
3 Geometric Airy Curve Flows	35
3.1 Equivalence of Geometric Airy Equations on \mathbb{R}^n and Vector Modified KdV Equation	35
3.2 Geometric Airy Curve Flows on \mathbb{S}^n	41
3.3 Geometric Airy Curve Flows on \mathbb{H}^n	44
4 Bäcklund Transformations	49
4.1 Bäcklund Transformations for $\frac{U}{K}$ -NLS and Schrödinger Flows	49
4.2 Bäcklund Transformations for Derivative $\frac{U}{K}$ -NLS	62
4.3 Bäcklund Transformations for vector mKdV and Geometric Airy Curve Flow on \mathbb{R}^n	70

5	Periodic Cauchy Problems	75
5.1	Periodic Cauchy Problems for Schrödinger Flow on S^2	75
5.2	Periodic Cauchy Problems for VFE on \mathbb{R}^3	79
5.3	Periodic Cauchy Problems for Geometric Airy Curve Flow on \mathbb{R}^2	88
6	Numerics	91
6.1	Geometric Algorithms	92
6.2	Numerical Experiments	96
	Bibliography	117
	Appendices	119
A	Appendix A	119

75
75
79
88

91
92
96

17

9
9

LIST OF FIGURES

	Page
6.1 Numerical solution $\gamma(x, t)$ of VFE where $\gamma(x, 0) = (2 \cos x, \sin x, 0)$. . .	98
6.2 Solution q of NLS corresponding to $\gamma(x, t)$ in Figure 6.1	99
6.3 Numerical solution $\gamma(x, t)$ of VFE where $\gamma(x, 0) = (\cos x, \sin x, \cos x)$	99
6.4 Solution q of NLS corresponding to $\gamma(x, t)$ in Figure 6.3	100
6.5 Numerical solution $\gamma(x, t)$ of VFE where $\gamma(x, 0) = (e^{0.1 \cos x}, \sin x, 0)$.	100
6.6 Solution q of NLS corresponding to $\gamma(x, t)$ in Figure 6.5	101
6.7 Apply BT to $\gamma(x, t) = (\cos x, \sin x, t)$	101
6.8 Apply BT to $\gamma(x, t)$ where $\gamma(x, 0) = (2 \cos x, \sin x, 0)$	102
6.9 $\gamma(x, t)$: the solution of Schrödinger flow on \mathbb{S}^2 with initial data $\gamma_0 =$ $(\cos x \cos(2x), \sin x \cos(2x), \sin(2x))$	107
6.10 Solution q of NLS corresponding to $\gamma(x, t)$ in Figure 6.9. The solid line represents the real part of q while the dashed one shows the imaginary part of q	107
6.11 $\gamma(x, t)$: the solution of Schrödinger flow on \mathbb{S}^2 with initial data $\gamma_0 =$ $(0.5 \sin(2x), \sin x, \sqrt{1 - 0.25 \sin^2(2x) - \sin^2 x})$	108
6.12 Solution q of NLS corresponding to $\gamma(x, t)$ in Figure 6.11. The solid line represents the real part of q while the dashed one shows the imaginary part of q	108
6.13 Apply BT to $\gamma(x, t) = (\cos x, \sin x, 0)$	109
6.14 Apply BT to $\gamma(x, t)$ in Fig. 6.9	110
6.15 Apply BT to $\gamma(x, t)$ in Fig. 6.11	110
6.16 Solution of Geometric Airy curve flow on \mathbb{R}^2 with initial data $\gamma_0 =$ $(2 \cos(2x), \sin(2x))$	112
6.17 Solution q of mKdV corresponding to $\gamma(x, t)$ in Figure 6.16. The solid line represents the real part of q and the dashed one shows the imagi- nary part of q	113
6.18 Solution of Geometric Airy curve flow on \mathbb{R}^2 with initial data $\gamma_0 =$ $(\cos x + 0.7 \sin x \cos x, \sin x + 0.7 \sin x \sin x)$	113
6.19 Solution q of mKdV corresponding to $\gamma(x, t)$ in Figure 6.18. The solid line shows the real part of q while the dashed one is the imaginary part of q	114
6.20 Solution of Geometric Airy curve flow on \mathbb{R}^2 with initial data $\gamma_0 =$ $(\sqrt{\sin(2x)} \cos x, \sqrt{\sin(2x)} \sin x)$	114

6.21	Solution q of mKdV corresponding to $\gamma(x, t)$ in Figure 6.20. The solid line shows the real part of q while the dashed one is the imaginary part of q .	115
6.22	Apply BT to $\gamma(x, t) = (\cos(x + \frac{t}{2}), \sin(x + \frac{t}{2}))$	115
6.23	Apply BT to the solution $\gamma(x, t)$ of geometric Airy curve flow on \mathbb{R}^2 with initial data $\gamma(x, 0) = (2 \cos x, \sin x)$	116
6.24	Apply BT to $\gamma(x, t)$ in Fig. 6.20	116

LIST OF TABLES

	Page
6.1 Errors with the true solution γ and the numerical solution γ_n	97
6.2 Conserved Quantities Error for VFE with initial data $\gamma_0 = (\cos x, \sin x, 0)$	103
6.3 Errors with the true solution γ and the numerical solution γ_n	105
6.4 Conserved Quantities Error for Schrödinger flow with initial data $\gamma_0 =$ $(\cos x, \sin x, 0)$	106
6.5 Errors with the true solution γ and the numerical solution γ_n	111
6.6 Conserved Quantity Errors for Airy curve Flow on \mathbb{R}^2 with initial data $\gamma_0 = (\cos x, \sin x)$	112
A.7 Conserved Quantity Errors for Schrödinger Flow on \mathbb{S}^2 with initial data $\gamma_0 = (0.5 \sin(2x), \sin x, \sqrt{1 - 0.25 \sin(2x)^2 - \sin^2 x})$	119
A.8 Conserved Quantity Errors for Schrödinger Flow on \mathbb{S}^2 with initial data $\gamma_0 = (\cos x \cos(2x), \sin x \cos(2x), \sin(2x))$	120
A.9 Conserved Quantities Error for VFE with initial data $\gamma_0 = (\sin x +$ $\cos x, \sin x, \cos x)$	121
A.10 Conserved Quantities Error for solution $\gamma(x, t)$ of VFE where $\gamma(x, 0) =$ $(\sin x \cos(2x), \sin x, \cos x)$	122
A.11 Conserved Quantity Errors for Airy curve Flow on \mathbb{R}^2 with initial data $\gamma_0 = (2 \cos x, \sin x)$	123
A.12 Conserved Quantity Errors for Airy curve Flow on \mathbb{R}^2 with initial data $\gamma_0 = (\cos x + 0.7 \sin x \cos x, \sin x + 0.7 \sin x \cos x)$	124