

The propagator of the attractive delta-Bose gas in one dimension

Sylvain Prolhac and Herbert Spohn

Citation: *Journal of Mathematical Physics* **52**, 122106 (2011); doi: 10.1063/1.3663431

View online: <http://dx.doi.org/10.1063/1.3663431>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/52/12?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Phase transition in the magnetic field in a Bose gas](#)

Low Temp. Phys. **36**, 283 (2010); 10.1063/1.3388802

[A heuristic approach to the weakly interacting Bose gas](#)

Am. J. Phys. **75**, 938 (2007); 10.1119/1.2766934

[Random matrix averages and the impenetrable Bose gas in Dirichlet and Neumann boundary conditions](#)

J. Math. Phys. **44**, 4157 (2003); 10.1063/1.1599954

[Graphical representation of the partition function of a one-dimensional \$\delta\$ -function Bose gas](#)

J. Math. Phys. **42**, 4883 (2001); 10.1063/1.1396836

[Thermodynamics of trapped interacting bosons in one dimension](#)

J. Math. Phys. **42**, 765 (2001); 10.1063/1.1331562



The propagator of the attractive delta-Bose gas in one dimension

Sylvain Prolhac^{a)} and Herbert Spohn^{b)}

Zentrum Mathematik and Physik Department, Technische Universität München, D-85747 Garching, Germany

(Received 23 September 2011; accepted 2 November 2011; published online 30 December 2011)

We consider the quantum δ -Bose gas on the infinite line. For repulsive interactions, Tracy and Widom have obtained an exact formula for the quantum propagator. In our contribution we explicitly perform its analytic continuation to attractive interactions. We also study the connection to the expansion of the propagator in terms of the Bethe ansatz eigenfunctions. Thereby we provide an independent proof of their completeness. © 2011 American Institute of Physics. [doi:10.1063/1.3663431]

I. INTRODUCTION

Quantum particles on the real line interacting through a δ -potential are governed by the Hamiltonian

$$H_\kappa = - \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} - 2\kappa \sum_{j < k}^n \delta(x_j - x_k). \quad (1.1)$$

The number of particles, n , is fixed throughout and $x = (x_1, \dots, x_n)$ denotes the positions of the particles. We will restrict ourselves to the bosonic subspace of symmetric wave functions. Equation (1.1) is the Hamiltonian of a quantum many-body system which can be analyzed through the Bethe ansatz. The repulsive interaction, $\kappa < 0$, has been studied in great detail and we refer to Refs. 1–7. The attractive case, $\kappa > 0$, has received less attention. One reason is that the structure of the Bethe equations is more complicated. On top, physical applications are not obviously in reach. In the recent years, there has been renewed interest. We have now available a detailed study of the eigenfunctions^{8–11} and, as argued by Calabrese and Caux,¹² applications to real materials are in sight. A further motivation comes from the one-dimensional Kadar-Parisi-Zhang (KPZ) equation.¹³ Its replica solution is given in terms of the propagator of the attractive δ -Bose gas,^{14,15} which can be used to obtain exact solutions for some special initial conditions.^{11,16–25}

In the KPZ context, and also in other cases, one is actually interested in the quantum propagator $\langle x | e^{-tH_\kappa} | y \rangle$, $t \geq 0$. In principle, e^{-tH_κ} can be expanded in a sum (integral) over eigenfunctions. But one might hope to have at disposal more concise expressions for the propagator. In the repulsive case, Tracy and Widom²⁶ carried out such a program. The resulting expression we refer to as TW formula, which will be discussed below, including its relation to the expansion in eigenfunctions. A natural issue is to extend such a program to the attractive case, which is the topic of our contribution.

By symmetry the propagator $\langle x | e^{-tH_\kappa} | y \rangle$ can be restricted to the domain $\Lambda = \{x | x_1 \leq \dots \leq x_n\} \subset \mathbb{R}^n$. Using the Bose symmetry, H_κ of (1.1) is then defined by

$$H_\kappa \psi(x_1, \dots, x_n) = - \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} \psi(x_1, \dots, x_n), \quad x \in \Lambda^\circ, \quad (1.2)$$

^{a)}Electronic mail: prolhac@ma.tum.de.

^{b)}Electronic mail: spohn@ma.tum.de.

with the boundary conditions

$$\left(\frac{\partial}{\partial x_{j+1}} - \frac{\partial}{\partial x_j} + \kappa \right) \psi(x_1, \dots, x_n) \Big|_{x_{j+1}=x_j} = 0, \quad (1.3)$$

where the limit $x_{j+1} = x_j$ is taken from the interior, Λ° , of Λ . The Hamiltonian H_κ is a self-adjoint operator and $\langle x | e^{-tH_\kappa} | y \rangle$ is continuous in $x, y \in \Lambda$. In particular,

$$\lim_{t \rightarrow 0} \langle x | e^{-tH_\kappa} | y \rangle = \frac{1}{n!} \prod_{j=1}^n \delta(x_j - y_j), \quad x, y \in \Lambda. \quad (1.4)$$

Throughout the paper $x, y \in \Lambda$, hence the position of the particles are ordered increasingly. As will be proved in the Appendix, $\langle x | e^{-tH_\kappa} | y \rangle$ is analytic in κ for otherwise fixed arguments. Thus, our strategy will be to analytically extend the TW formula, valid for $\kappa \leq 0$, to $\kappa > 0$. As written, the TW formula becomes singular at $\kappa = 0$. Therefore, the main task is to understand the structure of the analytic continuation in κ . As a result, we will arrive at various formulas for the propagator. One formula will be just the expansion in Bethe ansatz eigenfunctions, which thus implies their completeness.

The issue of completeness for the attractive δ -Bose gas on the line has been studied before. In his thesis, Oxford²⁷ proves completeness of the generalized eigenfunctions defined as bounded Bethe ansatz eigenfunctions. He uses functional analytic methods to construct the Hilbert space isometry from the generalized eigenfunctions and thereby the spectral representation of H_κ . A similar strategy is used by Babbitt and Thomas²⁸ for the ground state representation of the ferromagnetic Heisenberg model on \mathbb{Z} . Heckman and Opdam²⁹ exploit the fact that the δ -Bose gas turns up in the representation theory of graded Hecke algebras. (We are grateful to Balázs Pozsgay for pointing out this reference.) They have results for the case when the interaction strength is allowed to be pair dependent. But only for H_κ their expression simplifies and they arrive at a Plancherel formula, which is the completeness relation.

For the system on the line, studied here, the set of admissible wave numbers is known explicitly. For a bounded system, in particular with periodic boundary conditions, the discrete set of wave numbers are the solutions to the Bethe equations, a coupled system of n transcendental equations. Completeness becomes more difficult to establish and to our knowledge only for the repulsive case a completeness proof is available.³⁰

The article is organized as follows. In Sec. II, we recall the Tracy and Widom formula for the propagator in the repulsive case, and rewrite it in terms of Bethe eigenstates. In Sec. III, we summarize our main results on the propagator with attractive interactions. These results are proved in Sec. IV, by performing explicitly the analytic continuation to $\kappa > 0$. A further rewriting represents the propagator in terms of the known Bethe eigenstates. The special case of the propagator with all particles starting and ending at 0 is handled in Sec. V. In the Appendix, we prove that the (imaginary time) propagator is an analytic function of the coupling.

II. δ -BOSE GAS WITH REPULSIVE INTERACTION ($\kappa < 0$)

Let S_n be the set of all $n!$ permutations of the integers between 1 and n . In the following, we use the notations

$$\prod_{j < k}^n \equiv \prod_{1 \leq j < k \leq n} . \quad (2.1)$$

For $\kappa < 0$ the TW formula states (in the notation of Ref. 26 the strength of the potential is called $c = -\kappa$)

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \frac{1}{n!(2\pi)^n} \int_{\mathbb{R}^n} dq_1 \dots dq_n \\ &\sum_{\substack{\sigma \in S_n \\ \sigma(j) > \sigma(k)}} \prod_{\substack{j < k \\ \sigma(j) > \sigma(k)}} \frac{q_{\sigma(j)} - q_{\sigma(k)} - i\kappa}{q_{\sigma(j)} - q_{\sigma(k)} + i\kappa} \prod_{j=1}^n \left(e^{iq_{\sigma(j)}(x_{\sigma(j)} - y_j)} e^{-tq_j^2} \right). \end{aligned} \quad (2.2)$$

The connection to the eigenfunction expansion can be seen by symmetrizing over all permutations of the q_j . We introduce a new permutation $\tau \in S_n$ and replace q_j by $q_{\tau(j)}$. Replacing then σ by $\tau^{-1} \circ \sigma$, one finds

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \frac{1}{n!^2(2\pi)^n} \sum_{\sigma, \tau \in S_n} \int_{\mathbb{R}^n} dq_1 \dots dq_n \\ &\prod_{\substack{j, k=1 \\ \tau^{-1}(j) < \tau^{-1}(k) \\ \sigma^{-1}(j) > \sigma^{-1}(k)}}^n \frac{q_j - q_k + i\kappa}{q_j - q_k - i\kappa} \prod_{j=1}^n \left(e^{iq_j(x_{\sigma^{-1}(j)} - y_{\tau^{-1}(j)})} e^{-tq_j^2} \right). \end{aligned} \quad (2.3)$$

One notes the factorization

$$\begin{aligned} \prod_{\substack{j, k=1 \\ \tau^{-1}(j) < \tau^{-1}(k) \\ \sigma^{-1}(j) > \sigma^{-1}(k)}}^n \frac{q_j - q_k + i\kappa}{q_j - q_k - i\kappa} &= \prod_{\substack{j < k \\ \tau^{-1}(j) < \tau^{-1}(k) \\ \sigma^{-1}(j) > \sigma^{-1}(k)}}^n \frac{q_j - q_k + i\kappa}{q_j - q_k - i\kappa} \prod_{\substack{j < k \\ \tau^{-1}(j) > \tau^{-1}(k) \\ \sigma^{-1}(j) < \sigma^{-1}(k)}}^n \frac{q_j - q_k - i\kappa}{q_j - q_k + i\kappa} \\ &= \prod_{\substack{j < k \\ \sigma^{-1}(j) > \sigma^{-1}(k)}}^n \frac{q_j - q_k + i\kappa}{q_j - q_k - i\kappa} \prod_{\substack{j < k \\ \tau^{-1}(j) > \tau^{-1}(k)}}^n \frac{q_j - q_k - i\kappa}{q_j - q_k + i\kappa}. \end{aligned} \quad (2.4)$$

We introduce the Bethe eigenstates of the Hamiltonian (1.1) with repulsive interaction

$$\psi(x; q) = \frac{1}{n!} \sum_{\sigma \in S_n} \prod_{\substack{j < k \\ \sigma^{-1}(j) > \sigma^{-1}(k)}}^n \frac{q_j - q_k + i\kappa}{q_j - q_k - i\kappa} \prod_{j=1}^n e^{iq_j x_{\sigma^{-1}(j)}}, \quad (2.5)$$

with momenta $q = (q_1, \dots, q_n) \in \mathbb{R}^n$. The eigenstate $\psi(x; q)$ has energy

$$E(q) = \sum_{j=1}^n q_j^2. \quad (2.6)$$

In terms of Bethe eigenstates, Eqs. (2.3) and (2.4) rewrite as

$$\langle x | e^{-tH_\kappa} | y \rangle = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} dq_1 \dots dq_n \psi(x; q) \overline{\psi(y; q)} e^{-tE(q)}, \quad (2.7)$$

where $\overline{(\dots)}$ denotes complex conjugation. At $t = 0$, (2.7) reduces to the completeness relation for the Bethe eigenstates,

$$\mathbb{1} = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} dq_1 \dots dq_n |\psi(q)\rangle \langle \psi(q)|. \quad (2.8)$$

A proof of the orthonormality of the Bethe eigenstates can be found, e.g., in Appendix A of Ref. 11.

III. δ -BOSE GAS WITH ATTRACTIVE INTERACTION ($\kappa > 0$)

As shown in the Appendix, the propagator is an analytic function of the coupling κ for $t > 0$. By analytic continuation of (2.2) from $\kappa < 0$ to $\kappa > 0$, we will derive in Sec. III an exact expression for the propagator in the attractive case $\kappa > 0$.

Before stating the two main theorems, a few definitions are needed. We call $D_{n,M}$ the set of the M -tuples $\vec{n} = (n_1, \dots, n_M)$ such that $n_j \geq 1, j = 1, \dots, M$ and $n_1 + \dots + n_M = n$. For $\vec{n} \in D_{n,M}$, the clusters $\Omega_j(\vec{n}) \equiv \Omega_j, j = 1, \dots, M$ are defined by

$$\Omega_j = \{n_1 + \dots + n_{j-1} + 1, \dots, n_1 + \dots + n_j\}. \tag{3.1}$$

From the Bethe ansatz point of view, the clusters will correspond to bound states of the particles. The function $r_{\vec{n}} \equiv r$, acting on $\{1, \dots, n\}$, is defined by

$$r(a) = s \quad \text{for} \quad a = n_1 + \dots + n_{j-1} + s \in \Omega_j. \tag{3.2}$$

More visually, one has

$$\begin{array}{c|ccc|ccc|ccc|ccc} a & 1 & \dots & n_1 & n_1 + 1 & \dots & n_1 + n_2 & \dots & n - n_M + 1 & \dots & n \\ \text{Cluster} & \Omega_1 & & & \Omega_2 & & & \dots & \Omega_M & & \\ r(a) & 1 & \dots & n_1 & 1 & \dots & n_2 & \dots & 1 & \dots & n_M \end{array} \tag{3.3}$$

Finally, we call $S'_n(\vec{n})$ (respectively, $S''_n(\vec{n})$) the subset of S_n containing only the permutations σ (respectively, τ) such that for all $j = 1, \dots, M$ and $a, b \in \Omega_j$ with $a < b$ one has $\sigma^{-1}(a) < \sigma^{-1}(b)$ (respectively, $\tau^{-1}(a) > \tau^{-1}(b)$).

In the attractive case, the following expression for the propagator is proved in Sec. IV.

Theorem 1: For fixed \vec{n} , let $\mu_j, j = 1, \dots, M$, be arbitrary real numbers satisfying the constraint

$$-n_j < \mu_j \leq 0. \tag{3.4}$$

For $\kappa > 0$ and $x, y \in \Lambda$, one has

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{\kappa^{n-M}}{n! M! (2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \prod_{j=1}^M (n_j! (n_j - 1)!) \int_{\mathbb{R}^M} dq_1 \dots dq_M \\ &\sum_{\sigma \in S'_n(\vec{n})} \sum_{\tau \in S''_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} \left(e^{i(q_j + i\kappa(\mu_j + r(a) - 1))(x_{\sigma^{-1}(a)} - y_{\tau^{-1}(a)})} e^{-t(q_j + i\kappa(\mu_j + r(a) - 1))^2} \right) \\ &\times \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b) \\ \tau^{-1}(a) < \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa(\mu_j + r(a))) - (q_k + i\kappa(\mu_k + r(b))) + i\kappa}{(q_j + i\kappa(\mu_j + r(a))) - (q_k + i\kappa(\mu_k + r(b))) - i\kappa} \right). \end{aligned} \tag{3.5}$$

All the apparent poles in the integrand cancel except for simple poles at $q_j + i\kappa\mu_j = q_k + i\kappa(\mu_k + n_k)$ and $q_j + i\kappa(\mu_j + n_j) = q_k + i\kappa\mu_k, j < k$. The integrand vanishes at $q_j + i\kappa\mu_j = q_k + i\kappa\mu_k$ and $q_j + i\kappa(\mu_j + n_j) = q_k + i\kappa(\mu_k + n_k)$.

Compared to the TW formula (2.2), our result (3.5) is more complicated. Physically, the complication can be traced to the presence of bound states for the attractive case: the n particles are arranged in M clusters of size n_1, \dots, n_M , hence the extra summations over M and \vec{n} . Furthermore, Eq. (3.5) contains a summation over two permutations σ and τ instead of only one for the TW formula (2.2). In the special case $x = y = 0$ discussed in Sec. V, both summations over σ and τ can be eliminated.

For the attractive case, the propagator can also be written in terms of a summation over the eigenstates of the Hamiltonian (1.1). The Bethe eigenfunctions for attractive interaction are (see Eqs. (B.26) and (B.48) of Ref. 11; in Ref. 11 $r(a)$ is equal to $r(\sigma(a))$ with our notations, and Ω_j to

$\sigma^{-1}(\Omega_j)$

$$\begin{aligned} \psi(x; M, \vec{n}, q) = & \frac{\kappa^{\frac{n-M}{2}}}{\sqrt{n!}} \prod_{j=1}^M \sqrt{n_j!(n_j-1)!} \sum_{\sigma \in S_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} \left(e^{i(q_j + i\kappa(r(a) - \frac{n_j}{2} - \frac{1}{2}))x_{\sigma^{-1}(a)}} \right) \\ & \times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa(r(a) - \frac{n_j}{2})) - (q_k + i\kappa(r(b) - \frac{n_k}{2})) + i\kappa}{(q_j + i\kappa(r(a) - \frac{n_j}{2})) - (q_k + i\kappa(r(b) - \frac{n_k}{2})) - i\kappa} \right), \end{aligned} \tag{3.6}$$

with $M = 1, \dots, n, \vec{n} \in D_{n,M}$ and $q \in \mathbb{R}^M$. Eq. (3.6) is an eigenfunction of the Hamiltonian (1.1) with eigenvalue

$$E(M, \vec{n}, q) = \sum_{j=1}^M \left(n_j q_j^2 - \frac{\kappa^2}{12} (n_j^3 - n_j) \right). \tag{3.7}$$

The relation of the propagator with the Bethe eigenfunctions is stated as next theorem in terms of (3.6) and (3.7).

Theorem 2: For $\kappa > 0$ and $x, y \in \Lambda$, one has

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle = & \sum_{M=1}^n \frac{1}{M!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \int_{\mathbb{R}^M} dq_1 \dots dq_M \\ & \psi(x; M, \vec{n}, q) \overline{\psi(y; M, \vec{n}, q)} e^{-tE(M, \vec{n}, q)}. \end{aligned} \tag{3.8}$$

As in the case of repulsive interaction discussed in Sec. II, taking $t = 0$ yields the completeness relation for the Bethe eigenstates (3.6). Their orthonormality is proved in Appendix B of Ref. 11.

IV. ANALYTIC CONTINUATION FROM $\kappa < 0$ TO $\kappa > 0$

In this section, the TW formula (2.2) for the propagator is extended by analytic continuation to the attractive case $\kappa > 0$. Theorem 1 and Theorem 2 are proved.

A. Contribution of the residues

The contours of integration in (2.2) can be moved freely as long as the denominators $q_j - q_k - i\kappa$ keep a strictly positive imaginary part. In particular, if the integration is shifted to $q_j \in \mathbb{R} + i\lambda(n-j), j = 1, \dots, n$, with $\lambda > 0$, we obtain a formula valid for all κ such that $\Im(q_j - q_k - i\kappa) = (k-j)\lambda - \kappa > 0$ for $j < k$, i.e. for all $\kappa < \lambda$. One obtains

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle = & \frac{1}{n!(2\pi)^n} \prod_{a=1}^n \left(\int_{\mathbb{R} + i\lambda(n-a)} dq_a \right) \\ & \sum_{\sigma \in S_n} \prod_{\substack{a < b \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \frac{q_a - q_b + i\kappa}{q_a - q_b - i\kappa} \prod_{a=1}^n \left(e^{iq_a(x_{\sigma^{-1}(a)} - y_a)} e^{-tq_a^2} \right). \end{aligned} \tag{4.1}$$

In the following, we want to further move the contours of integration, but this time the contours will have to cross poles of the integrand, which will add several new terms resulting from the residues at

these poles, symbolically,

$$\begin{array}{c} \longrightarrow \\ \times \end{array} = \begin{array}{c} \textcircled{\times} \\ \longrightarrow \end{array} \tag{4.2}$$

If one denotes by $j \rightarrow k$ the action of taking the residue at $q_j = q_k + i\kappa r$ with $r \in \mathbb{Z}$, the terms, obtained after moving the contours of integration, correspond to collections of $j \rightarrow k$ such that, for each $\ell = 1, \dots, n$, $\ell \rightarrow \dots$ appears only once in the collection (since after taking the residue at $q_\ell = q_m + i\kappa r$, the integrand no longer contains q_ℓ). Each term thus corresponds to a forest (a set of trees), for example,

$$\{1 \rightarrow 2, 2 \rightarrow 7, 3 \rightarrow 5, 4 \rightarrow 7\} \Leftrightarrow \begin{array}{ccccc} 1 & & & & \\ \downarrow & & & & \\ 2 & & 4 & & 3 \\ & \searrow & \swarrow & & \downarrow \\ & 7 & & & 5 \end{array} \quad 6 \tag{4.3}$$

The particular trees obtained in this fashion depend on the order in which the contours are moved. Here, we choose to move first the contour for q_{n-1} in such a way that it crosses only the pole at $q_{n-1} = q_n + i\kappa$. Then, we move the contour for q_{n-2} in such a way that it crosses only the poles at $q_{n-2} = q_n + i\kappa$ and $q_{n-2} = q_{n-1} + i\kappa$ (in which case we still have an integration over both q_{n-1} and q_n), or only the pole at $q_{n-2} = q_n + 2i\kappa$ (in which case the residue at $q_{n-1} = q_n + i\kappa$ has been taken). We continue in this fashion until in the final step the contour for q_1 is moved.

In principle, after moving the contours, the propagator $\langle x|e^{-tH_\kappa}|y \rangle$ will be expressed as a sum over forests. In fact, it turns out that during this procedure there are many cancellations which remove all the forests which contain trees with ‘‘branches’’: in other words, only the forests with merely ‘‘branchless’’ trees (like $a, a \rightarrow b, a \rightarrow b \rightarrow c, a \rightarrow b \rightarrow c \rightarrow d, \dots$) remain after these cancellations. Instead of a sum over forests, we end up with a sum over partitions of $\{1, \dots, n\}$ (each element of the partition corresponding to one of the branchless trees of the forest).

In the context of the distribution of the leftmost particle in the asymmetric simple exclusion process, the procedure described here bears some similarity with the transformation from Theorem 3.1 to Theorem 3.2 in Ref. 31 where contours of integration are moved from small to large circles. A complication in our context is that we need a summation over all partitions of $\{1, \dots, n\}$ and not just over subsets of $\{1, \dots, n\}$. We expect that in the case of the full transition probability for the asymmetric exclusion process an expression with an integration over large circles would require summing over all partitions of $\{1, \dots, n\}$.

A proof of the previous statements is based on induction with respect to an integer ℓ such that all the contours for $q_{\ell+1}, \dots, q_{n-1}$ have already been moved.

We introduce a few notations. For a boolean condition c , $\mathbb{1}_{\{c\}}$ is defined to be equal to 1 if c is true and 0 otherwise. For $\vec{n} \in D_{n,M}$, the set $P_n(\vec{n})$ contains all the partitions $\vec{A} = \{A_1, \dots, A_M\}$ of $\{1, \dots, n\}$ with $|A_j| = n_j, j = 1, \dots, M$. The partition \vec{A} verifies $A_1 \cup \dots \cup A_M = \{1, \dots, n\}$ and for $j \neq k, A_j \cap A_k = \emptyset$. The partitions are not ordered, i.e., the partition $\vec{B} = \{A_{R(1)}, \dots, A_{R(M)}\}$ is considered to be the same element of $P_n(\vec{n})$ as \vec{A} for all $R \in S_M$. Each A_j is called a *cluster*, and will correspond to a bound state of particles in the Bethe ansatz point of view. For a partition \vec{A} , we define $d_{\vec{A}}(a), a = 1, \dots, n$, (abbreviated as $d(a)$ to lighten the notation) to be the rank of a in its cluster A_j , starting with rank 0 for the largest element of the cluster, rank 1 for the second largest, \dots , and rank $|A_j| - 1$ for the smallest element of A_j .

With these notation, the following lemma can be stated.

Lemma 1: Let ℓ be an integer between 0 and $n - 1$. For fixed $M = 1, \dots, n$, let $\epsilon_j, j = \ell + 1, \dots, M$ be distinct numbers with $0 \leq \epsilon_j < 1$. Then, for $0 < \kappa < \lambda$ one has

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{\kappa^{n-M}}{n!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \prod_{j=1}^M (n_j!(n_j - 1)!) \prod_{j=1}^{\ell} \left(\int_{\mathbb{R} + i\lambda(n-j)} dq_j \right) \\ &\times \prod_{j=\ell+1}^M \left(\int_{\mathbb{R} - i\kappa\epsilon_j} dq_j \right) \sum_{\vec{A} \in P_n(\vec{n})} \sum_{\sigma \in \mathcal{S}_n} \prod_{j=1}^{\ell} \mathbb{1}_{\{A_j = \{j\}\}} \prod_{j=1}^M \prod_{\substack{a, b \in A_j \\ a < b}} \mathbb{1}_{\{\sigma^{-1}(a) > \sigma^{-1}(b)\}} \\ &\times \prod_{j=1}^M \prod_{a \in A_j} \left(e^{i(q_j + i\kappa d(a))(x_{\sigma^{-1}(a)} - y_a)} e^{-t(q_j + i\kappa d(a))^2} \right) \\ &\times \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in A_j \\ b \in A_k \\ a < b \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) + i\kappa}{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) - i\kappa} \right). \end{aligned} \tag{4.4}$$

Proof: The constraint on the $\epsilon_j, j = \ell + 1, M$, implies that (4.4) is well defined since all the poles are at $q_j = q_k + i\delta$ with $\Im(q_j) \neq \Im(q_k) + \delta$.

For $\ell = n - 1$, the identity between the expressions (4.1) and (4.4) of $\langle x | e^{-tH_\kappa} | y \rangle$ is immediate. All the clusters must have size 1, and only $M = n$ contributes. Since the poles for q_n are at $q_n = q_j - i\kappa, j = 1, \dots, n - 1$, the contour for q_n can be moved freely from \mathbb{R} to $\mathbb{R} - i\kappa\epsilon_n$, provided $0 \leq \epsilon_n < 1$ and $\kappa > 0$.

We now proceed to prove the general identity by induction in ℓ : we assume that the expression (4.4) for $\langle x | e^{-tH_\kappa} | y \rangle$ is valid for $\ell \geq 1$ and will establish the expression with ℓ replaced by $\ell - 1$.

For given \vec{n} , we want to move the contour of integration for q_ℓ from $\mathbb{R} + i\lambda(n - \ell)$ to $\mathbb{R} - i\kappa\epsilon_\ell$ with $0 \leq \epsilon_\ell < 1$ and ϵ_ℓ different from all the other $\epsilon_k, k = \ell + 1, \dots, M$. In order to accomplish this, one needs to take into account the residues of the poles at $q_\ell = z$ for $-\kappa\epsilon_\ell < \Im(z) < \lambda(n - \ell)$. The only poles for q_ℓ are at $z = q_j - i\kappa, j = 1, \dots, \ell - 1$, and at $z = q_m + i\kappa(d(c) + 1), m = \ell + 1, \dots, M, c \in A_m$. In the first case, using $\kappa < \lambda$ and $j \leq \ell - 1$, one finds $\Im(z) > \lambda(n - \ell)$, which implies that these poles do not contribute when moving the contour for q_ℓ . In the second case, using $0 < \kappa < \lambda, 0 \leq \epsilon_m < 1$ and $\ell + d(c) + 1 \leq n$, one has $-\kappa\epsilon_\ell \leq 0 < \Im(z) < \lambda(n - \ell)$, which implies that all these poles contribute a residue (with a factor $-2i\pi$ corresponding to a clockwise contour integration).

Moving the contour for q_ℓ produces several terms: one term corresponding to the integration over $q_\ell \in \mathbb{R} - i\kappa\epsilon_\ell$, for which the integrand still depends on q_ℓ , and one term for each $c \in A_m, m = \ell + 1, \dots, M$, for which the residue at $q_\ell = q_m + i\kappa(d(c) + 1)$ has been taken. The latter term corresponds to merging the cluster $A_\ell = \{\ell\}$ and the cluster A_m . Assuming $\sigma^{-1}(\ell) > \sigma^{-1}(c)$ (otherwise, the pole vanishes), this term is equal to

$$\begin{aligned} &(-2i\pi) \frac{\kappa^{n-M}}{n!(2\pi)^M} \prod_{j=1}^M (n_j!(n_j - 1)!) \prod_{j=1}^{\ell-1} \left(\int_{\mathbb{R} + i\lambda(n-j)} dq_j \right) \prod_{j=\ell+1}^M \left(\int_{\mathbb{R} - i\kappa\epsilon_j} dq_j \right) \\ &\sum_{\vec{A} \in P_n(\vec{n})} \sum_{\sigma \in \mathcal{S}_n} \prod_{j=1}^{\ell} \mathbb{1}_{\{A_j = \{j\}\}} \prod_{j=1}^M \prod_{\substack{a, b \in A_j \\ a < b}} \mathbb{1}_{\{\sigma^{-1}(a) > \sigma^{-1}(b)\}} \\ &\times \left(e^{i(q_m + i\kappa(d(c)+1))(x_{\sigma^{-1}(\ell)} - y_\ell)} e^{-t(q_m + i\kappa(d(c)+1))^2} \right) \\ &\times \prod_{\substack{j=1 \\ j \neq \ell}}^M \prod_{a \in A_j} \left(e^{i(q_j + i\kappa d(a))(x_{\sigma^{-1}(a)} - y_a)} e^{-t(q_j + i\kappa d(a))^2} \right) \end{aligned}$$

$$\begin{aligned}
 & \times \prod_{\substack{j,k=1 \\ j \neq k \\ j,k \neq \ell}}^M \prod_{\substack{a \in A_j \\ b \in A_k \\ a < b \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) + i\kappa}{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) - i\kappa} \right) \\
 & \times \prod_{\substack{j,k=1 \\ j \neq k \\ j \neq \ell}}^M \prod_{\substack{a \in A_j \\ a < \ell \\ \sigma^{-1}(a) > \sigma^{-1}(\ell)}} \left(\frac{(q_j + i\kappa d(a)) - (q_m + i\kappa(d(c) + 1)) + i\kappa}{(q_j + i\kappa d(a)) - (q_m + i\kappa(d(c) + 1)) - i\kappa} \right) \\
 & \times \prod_{\substack{j,k=1 \\ j \neq k \\ k \neq \ell}}^M \prod_{\substack{b \in A_k \\ \ell < b \\ \sigma^{-1}(\ell) > \sigma^{-1}(b)}} \left(\frac{(q_m + i\kappa(d(c) + 1)) - (q_k + i\kappa d(b)) + i\kappa}{(q_m + i\kappa(d(c) + 1)) - (q_k + i\kappa d(b)) - i\kappa} \right) \\
 & \times \prod_{\substack{b \in A_m \\ b \neq c \\ \sigma^{-1}(\ell) > \sigma^{-1}(b)}} \left(\frac{(q_m + i\kappa(d(c) + 1)) - (q_m + i\kappa d(b)) + i\kappa}{(q_m + i\kappa(d(c) + 1)) - (q_m + i\kappa d(b)) - i\kappa} \right) \\
 & \times ((q_m + i\kappa(d(c) + 1)) - (q_m + i\kappa d(c)) + i\kappa). \tag{4.5}
 \end{aligned}$$

The last line of (4.5) contributes a factor $2i\kappa$ and the line before contributes

$$\prod_{\substack{b \in A_m \\ b \neq c \\ \sigma^{-1}(\ell) > \sigma^{-1}(b)}} \left(\frac{d(c) - d(b) + 2}{d(c) - d(b)} \right). \tag{4.6}$$

Let us first assume that $c = \min(A_m)$. Then, for all $b \in A_m, b \neq c$ one has $\sigma^{-1}(b) < \sigma^{-1}(c)$. Together with $\sigma^{-1}(c) < \sigma^{-1}(\ell)$, it implies that all the elements of the cluster A_m (except c) contribute in (4.6). This results in a factor $(n_m + 1)n_m/2$. Combined with $(-2i\pi)$ and $2i\kappa$, we obtain a factor $2\pi\kappa(n_k + 1)n_k$. The term with $c = \min(A_m)$ thus corresponds exactly to the term of (4.4) with ℓ replaced by $\ell - 1$ and the partition \vec{A} replaced by \vec{B} , obtained from \vec{A} by merging the cluster $\{\ell\}$ with A_m (after a renaming of the q_j, n_j, ϵ_j to $q_{j-1}, n_{j-1}, \epsilon_{j-1}$ for $\ell + 1 \leq j \leq M$).

It remains to show that for $c \neq \min(A_m)$, the residues cancel each other. Since the $\sigma^{-1}(b)$ are ordered in the same way as the $d(b)$ for $b \in A_m$, there exists a unique number $f \in A_m$ such that for $b \in A_m$, if $b \geq f$ then $\sigma^{-1}(b) < \sigma^{-1}(\ell)$, and if $b < f$ then $\sigma^{-1}(b) > \sigma^{-1}(\ell)$. Since $\sigma^{-1}(c) < \sigma^{-1}(\ell)$, one has necessarily $f \leq c$ (or equivalently $d(f) \geq d(c)$). Then, (4.6) rewrites

$$\prod_{\substack{b \in A_m \\ b \neq c \\ b \geq f}} \left(\frac{d(c) - d(b) + 2}{d(c) - d(b)} \right). \tag{4.7}$$

The rest of the argument depends on the relative values of $d(c)$ and $d(f)$. If $d(f) \geq d(c) + 2$, then, there exists $b \in A_m$ such that $b \geq f$ and $d(b) = d(c) + 2$, thus (4.7) is equal to zero. Since $d(f) \geq d(c)$, the only cases left are $d(f) = d(c) + 1$ and $d(f) = d(c)$, for which (4.7) rewrites, respectively,

$$\frac{d(c) - d(f) + 2}{d(c) - d(f)} \prod_{\substack{b \in A_m \\ b > c}} \left(\frac{d(c) - d(b) + 2}{d(c) - d(b)} \right) = -\frac{(d(c) + 1)(d(c) + 2)}{2}, \tag{4.8}$$

and

$$\prod_{\substack{b \in A_m \\ b > c}} \left(\frac{d(c) - d(b) + 2}{d(c) - d(b)} \right) = \frac{(d(c) + 1)(d(c) + 2)}{2}. \tag{4.9}$$

Let us call c' the element of A_m such that $d(c') = d(c) + 1$ (c' is the smallest element of A_m larger than c). One notes that the two previous cases are exchanged when replacing σ^{-1} by $\sigma^{-1} \circ \theta_{\ell, c'}$, with $\theta_{\ell, c'}$ the permutation exchanging ℓ and c' . Thus, summing over all permutations σ , the residues at $q_\ell = q_m + i\kappa(d(c) + 1)$ cancel. \square

Our construction achieves the proof of (4.4) for $0 < \kappa < \lambda$ and ℓ between 0 and $n - 1$. In particular, for $\ell = 0$, one has

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{\kappa^{n-M}}{n!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \prod_{j=1}^M (n_j!(n_j - 1)!) \\ &\times \prod_{j=1}^M \left(\int_{\mathbb{R} - i\kappa\epsilon_j} dq_j \right) \sum_{\vec{A} \in P_n(\vec{n})} \sum_{\sigma \in S_n} \prod_{j=1}^M \prod_{\substack{a, b \in A_j \\ a < b}} \mathbb{1}_{\{\sigma^{-1}(a) > \sigma^{-1}(b)\}} \\ &\times \prod_{j=1}^M \prod_{a \in A_j} \left(e^{i(q_j + i\kappa d(a))(x_{\sigma^{-1}(a)} - y_a)} e^{-t(q_j + i\kappa d(a))^2} \right) \\ &\times \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in A_j \\ b \in A_k \\ a < b \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) + i\kappa}{(q_j + i\kappa d(a)) - (q_k + i\kappa d(b)) - i\kappa} \right). \end{aligned} \tag{4.10}$$

This expression no longer depends on λ . Hence, it is valid in the entire range $\kappa > 0$.

B. Partitions and permutations

Exchanging A_j and A_k in (4.10) is the same as exchanging q_j and q_k , or ϵ_j and ϵ_k . Since the ϵ_j are arbitrary numbers satisfying a constraint ($0 \leq \epsilon_j < 1$ and all ϵ_j different) which is the same for all j , it is possible to add an extra sum over all permutations of the A_j , compensated by a global factor $1/M!$. This is equivalent to summing now over *ordered partitions* $\vec{A} = (A_1, \dots, A_n)$, such that for all $R \in S_M$ different from the identity permutation, the ordered partition $\vec{B} = (A_{R(1)}, \dots, A_{R(M)})$ is distinct from \vec{A} .

There exists a bijection between ordered partitions \vec{A} such that $|A_j| = n_j, j = 1, \dots, M$, and permutations $\tau \in S_n''(\vec{n})$ ($S_n''(\vec{n})$ is defined after Eq. (3.5)). By this bijection, the cluster A_j is equal to $\{\tau^{-1}(a), a \in \Omega_j(\vec{n})\} \equiv \tau^{-1}(\Omega_j)$ and one has $1 + d_{\vec{A}}(a) = r_{\vec{n}}(\tau(a))$, using the definitions (3.1) and (3.2). Equation (4.10) becomes

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{\kappa^{n-M}}{n!M!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \prod_{j=1}^M (n_j!(n_j - 1)!) \\ &\times \prod_{j=1}^M \left(\int_{\mathbb{R} - i\kappa\epsilon_j} dq_j \right) \sum_{\sigma \in S_n} \sum_{\tau \in S_n''(\vec{n})} \prod_{j=1}^M \prod_{\substack{a, b \in \tau^{-1}(\Omega_j) \\ a < b}} \mathbb{1}_{\{\sigma^{-1}(a) > \sigma^{-1}(b)\}} \end{aligned}$$

$$\begin{aligned} & \times \prod_{j=1}^M \prod_{a \in \tau^{-1}(\Omega_j)} \left(e^{i(q_j + i\kappa(r(\tau(a)) - 1))(x_{\sigma^{-1}(a)} - y_a)} e^{-t(q_j + i\kappa(r(\tau(a)) - 1))^2} \right) \\ & \times \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in \tau^{-1}(\Omega_j) \\ b \in \tau^{-1}(\Omega_k) \\ a < b \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa r(\tau(a))) - (q_k + i\kappa r(\tau(b))) + i\kappa}{(q_j + i\kappa r(\tau(a))) - (q_k + i\kappa r(\tau(b))) - i\kappa} \right). \end{aligned} \quad (4.11)$$

One can now replace everywhere a and b by $\tau^{-1}(a)$ and $\tau^{-1}(b)$. We also replace σ by $\tau^{-1} \circ \sigma$. Because of the definition of $S''_n(\vec{n})$, if $a, b \in \Omega_j$ with $\tau^{-1}(a) < \tau^{-1}(b)$ then $a > b$. This implies that the constraint with the $\mathbb{1}_{\{\dots\}}$ is equivalent to $\sigma \in S''_n(\vec{n})$ (defined after Eq. (3.5)). We obtain

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{\kappa^{n-M}}{n! M! (2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \prod_{j=1}^M (n_j! (n_j - 1)!) \prod_{j=1}^M \left(\int_{\mathbb{R} - i\kappa \epsilon_j} dq_j \right) \\ & \sum_{\sigma \in S''_n(\vec{n})} \sum_{\tau \in S''_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} \left(e^{i(q_j + i\kappa(r(a) - 1))(x_{\sigma^{-1}(a)} - y_{\tau^{-1}(a)})} e^{-t(q_j + i\kappa(r(a) - 1))^2} \right) \\ & \times \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b) \\ \tau^{-1}(a) < \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa} \right). \end{aligned} \quad (4.12)$$

C. Pole structure of the integrand and summation over Bethe eigenstates

One has the factorization

$$\begin{aligned} & \prod_{\substack{j, k = 1 \\ j \neq k}}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b) \\ \tau^{-1}(a) < \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa} \right) \\ &= \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b) \\ \tau^{-1}(a) < \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa} \right) \\ & \quad \times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) < \sigma^{-1}(b) \\ \tau^{-1}(a) > \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa} \right) \\ &= \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa} \right) \\ & \quad \times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \tau^{-1}(a) > \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa} \right). \end{aligned} \quad (4.13)$$

We introduce the functions

$$\begin{aligned} \varphi_\kappa(x; M, \vec{n}, q) &= \frac{\kappa^{\frac{n-M}{2}}}{\sqrt{n!}} \prod_{j=1}^M \sqrt{n_j!(n_j - 1)!} \sum_{\sigma \in S'_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} (e^{i(q_j + i\kappa(r(a)-1))x_{\sigma^{-1}(a)}}) \\ &\times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa} \right), \end{aligned} \quad (4.14)$$

and

$$\begin{aligned} \tilde{\varphi}_\kappa(y; M, \vec{n}, q) &= \frac{\kappa^{\frac{n-M}{2}}}{\sqrt{n!}} \prod_{j=1}^M \sqrt{n_j!(n_j - 1)!} \sum_{\tau \in S''_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} (e^{-i(q_j + i\kappa(r(a)-1))y_{\tau^{-1}(a)}}) \\ &\times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \tau^{-1}(a) > \tau^{-1}(b)}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) - i\kappa}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa} \right). \end{aligned} \quad (4.15)$$

One notes that $\tau \in S''_n(\vec{n})$ is equivalent to $R(\vec{n}) \circ \tau \in S'_n(\vec{n})$ with $R(\vec{n})$ the permutation that exchanges $a, b \in \Omega_j, j = 1, \dots, M$, if $r(a) + r(b) = n_j$. This implies

$$\tilde{\varphi}_\kappa(y; M, \vec{n}, q) = \overline{\varphi_\kappa(y; M, \vec{n}, q - i\kappa\vec{n} + i\kappa)}, \quad (4.16)$$

where $\overline{(\dots)}$ denotes complex conjugation. Thus, one can write

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{1}{M!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \left(\int_{\mathbb{R} - i\kappa\epsilon_j} dq_j \right) \\ &\prod_{j=1}^M \prod_{a \in \Omega_j} \left(e^{-t(q_j + i\kappa(r(a)-1))^2} \right) \varphi_\kappa(x; M, \vec{n}, q) \overline{\varphi_\kappa(y; M, \vec{n}, q - i\kappa\vec{n} + i\kappa)}. \end{aligned} \quad (4.17)$$

At this point, one wants to move again the contours of integration so that the exact Bethe eigenfunctions appear. One then needs to know precisely the location of the poles of the integrand in (4.17). To achieve this, we prove the following lemma:

Lemma 2: The expression

$$\varphi_\kappa(x; M, \vec{n}, q) \prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left(\frac{q_j - q_k + i\kappa(r - s - 1)}{q_j - q_k + i\kappa(r - s)} \right), \quad (4.18)$$

with $\varphi_\kappa(x; M, \vec{n}, q)$ defined in Eq. (4.14), is holomorphic as function of the q_j 's in the whole complex plane.

Proof: One defines $\chi(x; M, \vec{n}, q)$, equal to (4.18) up to a global normalization independent of q , by

$$\begin{aligned} \chi(x; M, \vec{n}, q) &= \sum_{\sigma \in S'_n(\vec{n})} \prod_{j=1}^M \prod_{a \in \Omega_j} (e^{i(q_j + i\kappa r(a) - i\kappa)x_{\sigma^{-1}(a)}}) \\ &\times \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k}} \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa \operatorname{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b))} \right). \end{aligned} \tag{4.19}$$

Noting that the product

$$\prod_{\substack{a, b \in \Omega_j \\ a < b}} \left(\frac{(q_j + i\kappa r(a)) - (q_j + i\kappa r(b)) + i\kappa \operatorname{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{(q_j + i\kappa r(a)) - (q_j + i\kappa r(b))} \right) \tag{4.20}$$

is nonzero for all $j = 1, \dots, M$ if and only if the permutation σ is an element of $S'_n(\vec{n})$, one can instead sum over all permutations $\sigma \in S_n$. One has

$$\begin{aligned} \chi(x; M, \vec{n}, q) &= \prod_{j=1}^M n_j! \sum_{\sigma \in S_n} \prod_{j=1}^M \prod_{a \in \Omega_j} (e^{i(q_j + i\kappa r(a) - i\kappa)x_{\sigma^{-1}(a)}}) \\ &\times \prod_{j < k}^M \prod_{a < b}^n \left(\frac{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b)) + i\kappa \operatorname{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{(q_j + i\kappa r(a)) - (q_k + i\kappa r(b))} \right). \end{aligned} \tag{4.21}$$

Then, one defines $\tilde{\chi}(x; M, \vec{n}, \xi)$ by replacing everywhere in $\chi(x; M, \vec{n}, q)$ the variables $q_j + i\kappa(r(a) - 1), j = 1, \dots, M, a \in \Omega_j$ by ξ_a with the result

$$\begin{aligned} \tilde{\chi}(x; M, \vec{n}, \xi) &= \prod_{j=1}^M n_j! \sum_{\sigma \in S_n} \prod_{a=1}^n (e^{i\xi_a x_{\sigma^{-1}(a)}}) \\ &\times \prod_{a < b}^n \left(\frac{\xi_a - \xi_b + i\kappa \operatorname{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{\xi_a - \xi_b} \right). \end{aligned} \tag{4.22}$$

If $\tilde{\chi}(x; M, \vec{n}, \xi)$ is analytic in the ξ_a , then $\chi(x; M, \vec{n}, q)$ will also be analytic in the q_j . One notes that $\tilde{\chi}(x; M, \vec{n}, \xi)$ has only simple poles. The term with the permutation $\sigma \in S_n$ of the residue at ξ_c

= ξ_d with $c < d$ is given by

$$\begin{aligned}
 & \left(\prod_{j=1}^M n_j! \right) e^{i\xi_d(x_{\sigma^{-1}(c)} + x_{\sigma^{-1}(d)})} \prod_{\substack{a=1 \\ a \neq c, d}}^n (e^{i\xi_a x_{\sigma^{-1}(a)}}) \\
 & \times i\kappa \operatorname{sign}(\sigma^{-1}(c) - \sigma^{-1}(d)) \\
 & \times \prod_{\substack{a < b \\ a \neq c, d \\ b \neq c, d}}^n \left(\frac{\xi_a - \xi_b + i\kappa \operatorname{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{\xi_a - \xi_b} \right) \\
 & \times \prod_{\substack{b=1 \\ b \neq c, d}}^n \left(\frac{\xi_d - \xi_b + i\kappa \operatorname{sign}(\sigma^{-1}(c) - \sigma^{-1}(b))}{\xi_d - \xi_b} \right) \\
 & \times \prod_{\substack{b=1 \\ b \neq c, d}}^n \left(\frac{\xi_d - \xi_b + i\kappa \operatorname{sign}(\sigma^{-1}(d) - \sigma^{-1}(b))}{\xi_d - \xi_b} \right). \tag{4.23}
 \end{aligned}$$

This term cancels with the term for which $\sigma^{-1}(a)$ and $\sigma^{-1}(b)$ are exchanged. Thus, $\tilde{\chi}(x; M, \vec{n}, \xi)$ and $\chi(x; M, \vec{n}, q)$ are analytic functions in the ξ_a and in the q_j . This finishes the proof of the lemma. \square

Then, since

$$\begin{aligned}
 & \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left(\frac{q_j - q_k + i\kappa(r - s - 1)}{q_j - q_k + i\kappa(r - s)} \frac{q_j - q_k + i\kappa(n_j - n_k - r + s + 1)}{q_j - q_k + i\kappa(n_j - n_k - r + s)} \right) \\
 & = \frac{(q_j - q_k - i\kappa n_k)(q_j - q_k + i\kappa n_j)}{(q_j - q_k)(q_j - q_k + i\kappa(n_j - n_k))}, \tag{4.24}
 \end{aligned}$$

the integrand in (4.17) is analytic in the q_j except for simple poles at $q_j = q_k + i\kappa n_k$ and $q_k = q_j + i\kappa n_j$ for $1 \leq j < k \leq M$. At this point, it is possible to set $\epsilon_j = 0$ for all $j = 1, \dots, M$ in (4.17) since the poles at $q_j = q_k$ cancel. Then, the contours of integration can be moved again, from $q_j \in \mathbb{R}$ to $q_j \in \mathbb{R} + i\mu_j$. Before moving the contours, one has (with $\kappa > 0$ and $j < k$)

$$\Im(q_j - q_k - i\kappa n_k) = -\kappa n_k < 0 \quad \text{and} \quad \Im(q_k - q_j - i\kappa n_j) = -\kappa n_j < 0. \tag{4.25}$$

After moving the contours, these inequalities must still be valid. One notes that if the μ_j 's satisfy the constraint

$$-n_j < \mu_j \leq 0, \tag{4.26}$$

both inequalities are satisfied. Shifting the contours and making then the change of variables $q_j \rightarrow q_j + i\kappa \mu_j$ in (4.12) leads to the result (3.5) of Theorem 1, while shifting the contours and making the change of variables in (4.17) leads to

$$\begin{aligned}
 \langle x | e^{-tH_\kappa} | y \rangle & = \sum_{M=1}^n \frac{1}{M!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \int_{\mathbb{R}^M} dq_1 \dots dq_M \prod_{j=1}^M \prod_{a \in \Omega_j} \left(e^{-t(q_j + i\kappa(\mu_j + r(a)-1))^2} \right) \\
 & \times \overline{\varphi_\kappa(x; M, \vec{n}, q + i\kappa \vec{\mu})} \varphi_\kappa(y; M, \vec{n}, q + i\kappa \vec{\mu} + i\kappa \vec{n} - i\kappa). \tag{4.27}
 \end{aligned}$$

One notes that

$$\begin{aligned} \sum_{a \in \Omega_j} (q_j + i\kappa(\mu_j + r(a) - 1))^2 &= n_j q_j^2 + 2i\kappa q_j \sum_{r=1}^{n_j} (\mu_j + r(a) - 1) - \kappa^2 \sum_{r=1}^{n_j} (\mu_j + r - 1)^2 \\ &= n_j q_j^2 + 2i\kappa q_j n_j \left(\mu_j + \frac{(n_j - 1)}{2} \right) - \frac{\kappa^2}{12} (n_j^3 - n_j) - \kappa^2 n_j \left(\mu_j + \frac{(n_j - 1)}{2} \right)^2. \end{aligned} \quad (4.28)$$

The choice $\mu_j = -(n_j - 1)/2$ is thus the only one such that the argument of the exponential in (4.27) is a real number. It is also the only choice such that $\varphi_\kappa(x; M, \vec{n}, q + i\kappa\vec{\mu})$ and $\overline{\varphi_\kappa(y; M, \vec{n}, q + i\kappa\vec{\mu} + i\kappa\vec{n} - i\kappa)}$ are complex conjugates of each other. Introducing the Bethe eigenfunctions¹¹

$$\psi(x; M, \vec{n}, q) = \varphi_\kappa \left(x; M, \vec{n}, q - \frac{i\kappa}{2} \vec{n} + \frac{i\kappa}{2} \right), \quad (4.29)$$

one finds for $\mu_j = -(n_j - 1)/2$

$$\begin{aligned} \langle x | e^{-tH_\kappa} | y \rangle &= \sum_{M=1}^n \frac{1}{M!(2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \int_{\mathbb{R}^M} dq_1 \dots dq_M \\ \psi(x; M, \vec{n}, q) \overline{\psi(y; M, \vec{n}, q)} &= \prod_{j=1}^M e^{-n_j q_j^2 + \frac{i\kappa^2}{12} (n_j^3 - n_j)}. \end{aligned} \quad (4.30)$$

This is the result (3.8) of Theorem 2.

V. THE SPECIAL CASE $x = y = 0$

In the special case $x = y = 0$, Eq. (4.30) simplifies. One has

$$\begin{aligned} \sum_{\sigma \in S'_n(\vec{n})} \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k \\ \sigma^{-1}(a) > \sigma^{-1}(b)}} \left(\frac{(q_j + i\kappa(\mu_j + r(a))) - (q_k + i\kappa(\mu_k + r(b))) + i\kappa}{(q_j + i\kappa(\mu_j + r(a))) - (q_k + i\kappa(\mu_k + r(b))) - i\kappa} \right) \\ = \frac{\sum_{\sigma \in S'_n(\vec{n})} \prod_{j < k}^M \prod_{\substack{a \in \Omega_j \\ b \in \Omega_k}} \left((q_j - q_k) + i\kappa(\mu_j - \mu_k + r(a) - r(b) + \text{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))) \right)}{\prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left((q_j - q_k) + i\kappa(\mu_j - \mu_k + r - s - 1) \right)}. \end{aligned} \quad (5.1)$$

As in Sec. IV, one can sum instead over all permutations $\sigma \in S_n$ in case the product in the numerator becomes a product over all $a < b$: the permutations which do not belong to $S'_n(\vec{n})$ give zero contribution. We use the notation $\xi_a = q_j + i\kappa(\mu_j + r(a))$ for $a \in \Omega_j$. Since the signature of the permutation σ can be written

$$\text{sign}(\sigma) = \prod_{a < b} \frac{\xi_a - \xi_b + i\kappa \text{sign}(\sigma^{-1}(a) - \sigma^{-1}(b))}{\xi_{\sigma(a)} - \xi_{\sigma(b)} + i\kappa \text{sign}(a - b)}, \quad (5.2)$$

Eq. (5.1) rewrites as

$$(-i\kappa)^{\frac{n_j(n_j-1)}{2}} \left(\prod_{j=1}^M \prod_{r=1}^{n_j} r! \right) \frac{\sum_{\sigma \in S_n} \text{sign}(\sigma) \prod_{a < b} \left(\xi_{\sigma(a)} - \xi_{\sigma(b)} + i\kappa \text{sign}(a - b) \right)}{\prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left((q_j - q_k) + i\kappa(\mu_j - \mu_k + r - s - 1) \right)}. \quad (5.3)$$

One has (e.g., see Lemma 1 of Ref. 19 for a proof)

Lemma 3: Let $f(a, b)$ be arbitrary complex coefficients. Then

$$\sum_{\sigma \in \mathcal{S}_n} \text{sign}(\sigma) \prod_{a < b}^n (\xi_{\sigma(a)} - \xi_{\sigma(b)} + f(a, b)) = n! \prod_{a < b}^n (\xi_a - \xi_b). \quad (5.4)$$

Using this lemma, (5.3) rewrites

$$\begin{aligned} & n! (-i\kappa)^{-\frac{n_j(n_j-1)}{2}} \left(\prod_{j=1}^M \prod_{r=1}^{n_j} \frac{1}{r!} \right) \frac{\prod_{a < b}^n (\xi_a - \xi_b)}{\prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} ((q_j - q_k) + i\kappa(\mu_j - \mu_k + r - s - 1))} \\ &= n! \left(\prod_{j=1}^M \frac{1}{n_j!} \right) \prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left(\frac{(q_j - q_k) + i\kappa(\mu_j - \mu_k + r - s)}{(q_j - q_k) + i\kappa(\mu_j - \mu_k + r - s - 1)} \right). \end{aligned} \quad (5.5)$$

Then using (see Eqs. (B.55)–(B.58) and (34) of Ref. 11)

$$\begin{aligned} & \prod_{j < k}^M \prod_{r=1}^{n_j} \prod_{s=1}^{n_k} \left| \frac{(q_j - q_k) + i\kappa(-\frac{n_j}{2} + \frac{n_k}{2} + r - s)}{(q_j - q_k) + i\kappa(-\frac{n_j}{2} + \frac{n_k}{2} + r - s - 1)} \right|^2 \\ &= \prod_{j < k}^M \left| \frac{(q_j - q_k) - \frac{i\kappa}{2}(n_j - n_k)}{(q_j - q_k) - \frac{i\kappa}{2}(n_j + n_k)} \right|^2 \\ &= \det \left(\frac{i\kappa n_j}{(q_j - q_k) + \frac{i\kappa}{2}(n_j + n_k)} \right)_{j,k=1,\dots,M}, \end{aligned} \quad (5.6)$$

one finds

$$\begin{aligned} \langle 0 | e^{-tH_\kappa} | 0 \rangle &= \sum_{M=1}^n \frac{n! \kappa^n}{M! (2\pi)^M} \sum_{\vec{n} \in D_{n,M}} \int_{\mathbb{R}^M} dq_1 \dots dq_M \\ & \det \left(\frac{e^{-n_j q_j^2 + \frac{i\kappa^2}{12}(n_j^3 - n_j)}}{-i(q_j - q_k) + \frac{\kappa}{2}(n_j + n_k)} \right)_{j,k=1,\dots,M}. \end{aligned} \quad (5.7)$$

In the replica computations for the KPZ equation with sharp wedge initial data, Eq. (5.7) can be used to arrive directly at the generating function of the height fluctuations, see Eqs. (2.17)–(2.19) of Ref. 19.

VI. CONCLUSIONS

Exact formulas for the transition probability of the one-dimensional asymmetric simple exclusion process, a non-equilibrium exactly solvable model, have been derived a few years ago.^{31–33} Subsequently, the method was adapted to obtain an exact formula for the propagator of the quantum δ -Bose gas with repulsive interaction.²⁶ Here, we analytically continued this formula to the case of attractive interaction.

An advantage of our approach, compared to the usual Bethe ansatz, is that the question of the completeness of the Bethe eigenfunctions can be completely avoided. In fact, such kind of exact expressions for the propagator can be used to *prove* the completeness of the Bethe ansatz, at least on the infinite line. In principle, it might also be possible to use the same kind of approach for the case of periodic boundary conditions, see, e.g., Refs. 33 and 34.

The exact expression for the propagator of the repulsive δ -Bose gas, derived in Ref. 26 and used here, bears some formal similarity with the coordinate Bethe ansatz, which is the original ansatz introduced by Bethe to diagonalize the Hamiltonian of Heisenberg spin chain. Since then, other

descriptions of eigenstates have been developed, in particular algebraic Bethe ansatz, which makes clearer the structures underlying the quantum integrability of such type of models. It would be of interest to understand whether it is possible to write down the propagator using an approach closer to the algebraic Bethe ansatz.

ACKNOWLEDGMENTS

We thank Tomohiro Sasamoto for most instructive discussions. S.P. acknowledges the support through a Deutsche Forschungsgemeinschaft (DFG) research project (SP181-24).

APPENDIX: ANALYTICITY IN THE COUPLING OF THE PROPAGATOR OF THE δ -BOSE GAS

In the case of a standard Schrödinger operator of the form $-\Delta + \lambda V$ one can use Kato's theory to establish that $e^{-t(-\Delta + \lambda V)}$ is analytic in λ . The δ -potential corresponds to a boundary condition and we are not aware of a functional analytic argument for the holomorphic dependence on κ . Instead, we will use the Feynman-Kac representation.

Proposition 1: For fixed $x, y \in \mathbb{R}^n$, $t > 0$, the function $\kappa \mapsto \langle x | e^{-tH_\kappa} | y \rangle$ is holomorphic on \mathbb{C} .

Proof: By the Feynman-Kac formula one has the representation

$$\langle x | e^{-tH_\kappa} | y \rangle = \mathbb{E}_{x,y} (e^{2\kappa X(t)}) p_t(x - y). \quad (\text{A1})$$

Here $p_t(x - y)$ is the Brownian motion transition kernel. The expectation $\mathbb{E}_{x,y}$ is over the standard Brownian bridge, $b(t)$, starting at x and ending at y at time t . Let $L_{j,k}(t)$, $j < k$, be the local time at 0 for $\{b_i(s) - b_j(s), 0 \leq s \leq t\}$, i.e.,

$$L_{j,k}(t) = \int_0^t ds \delta(b_j(s) - b_k(s)). \quad (\text{A2})$$

Then

$$X(t) = \sum_{j < k}^n L_{j,k}(t). \quad (\text{A3})$$

Since $X(t) \geq 0$, we bound as

$$\mathbb{E}_{x,y} (e^{2\kappa X(t)}) \leq \mathbb{E}_{x,y} (e^{2|\kappa|X(t)}) \leq \sum_{j < k}^n \mathbb{E}_{x,y} (e^{|\kappa|n^2 L_{j,k}(t)}). \quad (\text{A4})$$

The second inequality follows from the convexity of the exponential function. $L_{j,k}(t)$ is the local time at 0 for a one-dimensional Brownian bridge starting at $x_j - x_k$ and ending at $y_j - y_k$ at time t . The distribution of $L_{j,k}(t)$ has a Gaussian decay at infinity. Hence for any $|\kappa|$ the right-hand side in (A4) is bounded. This proves analyticity on \mathbb{C} . \square

¹ E. H. Lieb and W. Liniger, "Exact analysis of an interacting Bose gas. I. The general solution and the ground state," *Phys. Rev.* **130**, 1605–1616 (1963).

² E. H. Lieb, "Exact analysis of an interacting Bose gas. II. The excitation spectrum," *Phys. Rev.* **130**, 1616–1624 (1963).

³ C. N. Yang and C. P. Yang, "Thermodynamics of a one-dimensional system of bosons with repulsive delta-function interaction," *J. Math. Phys.* **10**, 1115–1122 (1969).

⁴ M. Gaudin, "Bose gas in one dimension. I. The closure property of the scattering wavefunctions," *J. Math. Phys.* **12**, 1674–1676 (1971).

⁵ M. Gaudin, "Bose gas in one dimension. II. Orthogonality of the scattering states," *J. Math. Phys.* **12**, 1677–1680 (1971).

⁶ M. Gaudin, "Boundary energy of a Bose gas in one dimension," *Phys. Rev. A* **4**, 386–394 (1971).

⁷ M. Gaudin, *La Fonction d'Onde de Bethe* (Masson, Paris, 1983).

⁸ J. B. McGuire, "Study of exactly soluble one-dimensional N-body problems," *J. Math. Phys.* **5**, 622–636 (1964).

⁹ A. N. Kirillov and V. E. Korepin, "Norms of bound states," *J. Math. Sci.* **40**, 13–21 (1988).

¹⁰ P. Calabrese and J. S. Caux, "Dynamics of the attractive 1D Bose gas, analytical treatment from integrability," *J. Stat. Mech.* P08032 (2007).

- ¹¹ V. Dotsenko, "Replica Bethe ansatz derivation of the Tracy-Widom distribution of the free energy fluctuations in one-dimensional directed polymers," *J. Stat. Mech.* P07010 (2010).
- ¹² P. Calabrese and J. S. Caux, "Correlation functions of the one-dimensional attractive Bose gas," *Phys. Rev. Lett.* **98**, 150403 (2007).
- ¹³ M. Kardar, G. Parisi, and Y.-C. Zhang, "Dynamic scaling of growing interfaces," *Phys. Rev. Lett.* **56**, 889–892 (1986).
- ¹⁴ M. Kardar, "Replica Bethe ansatz studies of two-dimensional interfaces with quenched random impurities," *Nucl. Phys. B* **290**, 582–602 (1987).
- ¹⁵ J. P. Bouchaud and H. Orland, "On the Bethe ansatz for random directed polymers," *J. Stat. Phys.* **61**, 877–884 (1990).
- ¹⁶ V. Dotsenko and B. Klumov, "Bethe ansatz solution for one-dimensional directed polymers in random media," *J. Stat. Mech.* P03022 (2010).
- ¹⁷ P. Calabrese, P. Le Doussal, and A. Rosso, "Free-energy distribution of the directed polymer at high temperature," *Europhys. Lett.* **90**, 20002 (2010).
- ¹⁸ V. Dotsenko, "Bethe ansatz derivation of the Tracy-Widom distribution for one-dimensional directed polymers," *Europhys. Lett.* **90**, 20003 (2010).
- ¹⁹ S. Prolhac and H. Spohn, "Two-point generating function of the free energy for a directed polymer in a random medium," *J. Stat. Mech.* P01031 (2011).
- ²⁰ S. Prolhac and H. Spohn, "The one-dimensional KPZ equation and the airy process," *J. Stat. Mech.* P03020 (2011).
- ²¹ S. Prolhac and H. Spohn, "Height distribution of the Kardar-Parisi-Zhang equation with sharp-wedge initial condition: Numerical evaluations," *Phys. Rev. E* **84**, 011119 (2011).
- ²² P. Calabrese and P. Le Doussal, "Exact solution for the Kardar-Parisi-Zhang equation with flat initial conditions," *Phys. Rev. Lett.* **106**, 250603 (2011).
- ²³ T. Imamura and T. Sasamoto, "Replica approach to the KPZ equation with the half Brownian motion initial condition," *J. Phys. A: Math. Theor.* **44**, 385001 (2011).
- ²⁴ I. Corwin and J. Quastel, e-print arXiv:1103.3422.
- ²⁵ I. Corwin, e-print arXiv:1106.1596.
- ²⁶ C. A. Tracy and H. Widom, "The dynamics of the one-dimensional delta-function Bose gas," *J. Phys. A: Math. Theor.* **41**, 485204 (2008).
- ²⁷ S. C. Oxford, "The hamiltonian of the quantized non-linear Schrödinger equation," Ph.D. dissertation (University of California, Los Angeles (UCLA), 1979).
- ²⁸ D. Babbitt and L. Thomas, "Ground state representation of the infinite one-dimensional Heisenberg ferromagnet," *Comm. Math. Phys.* **54**, 255–278 (1977).
- ²⁹ G. J. Heckman and E. M. Opdam, "Yang's system of particles and Hecke algebras," *Ann. Math.* **145**, 139–173 (1997).
- ³⁰ T. Dorlas, "Orthogonality and completeness of the Bethe ansatz eigenstates of the nonlinear Schrödinger model," *Commun. Math. Phys.* **154**, 347–376 (1993).
- ³¹ C. A. Tracy and H. Widom, "Integral formulas for the asymmetric simple exclusion process," *Commun. Math. Phys.* **279**, 815–844 (2008).
- ³² G. M. Schütz, "Exact solution of the master equation for the asymmetric exclusion process," *J. Stat. Phys.* **88**, 427–445 (1997).
- ³³ V. B. Priezzhev, "Exact nonstationary probabilities in the asymmetric exclusion process on a ring," *Phys. Rev. Lett.* **91**, 050601 (2003).
- ³⁴ A. M. Povolotsky and V. B. Priezzhev, "Determinant solution for the totally asymmetric exclusion process with parallel update: II. Ring geometry," *J. Stat. Mech.* P08018 (2007).